

# **The Role of the Physical Environment in the Hospital of the 21<sup>st</sup> Century: A Once-in-a-Lifetime Opportunity**

Roger Ulrich\*, Xiaobo Quan, Center for Health Systems and Design, College of Architecture, Texas A&M University

Craig Zimring\*, Anjali Joseph, Ruchi Choudhary, College of Architecture, Georgia Institute of Technology

\* Co-principal investigators and corresponding authors: Roger Ulrich, Ph.D.: [ulrich@archone.tamu.edu](mailto:ulrich@archone.tamu.edu); Craig Zimring, Ph.D.: [craig.zimring@arch.gatech.edu](mailto:craig.zimring@arch.gatech.edu).

Report to The Center for Health Design for the *Designing the 21<sup>st</sup> Century Hospital Project*. This project is funded by the Robert Wood Johnson Foundation.

September 2004

## **The Role of the Physical Environment in the Hospital of the 21<sup>st</sup> Century: A Once-in-a-Lifetime Opportunity**

A visit to a U.S. hospital is dangerous and stressful for patients, families and staff members. Medical errors and hospital-acquired infections are among the leading causes of death in the United States, each killing more Americans than AIDS, breast cancer, or automobile accidents (Institute of Medicine, 2000; 2001). According to the Institute of Medicine in its landmark *Quality Chasm* report: “*The frustration levels of both patients and clinicians have probably never been higher. Yet the problems remain. Health care today harms too frequently and routinely fails to deliver its potential benefits*” (IOM, 2001). Problems with U.S. health care not only influence patients; they impact staff. Registered nurses have a turnover rate averaging 20 percent (Joint Commission on Accreditation of Healthcare Organizations, 2002).

At the same time, the United States is facing one of the largest hospital building booms in US history. As a result of a confluence of the need to replace aging 1970s hospitals, population shifts in the United States, the graying of the baby boom generation, and the introduction of new technologies, the United States will spend more than \$16 billion for hospital construction in 2004, and this will rise to more than \$20 billion per year by the end of the decade (Babwin, 2002). These hospitals will remain in place for decades.

This once-in-lifetime construction program provides an opportunity to rethink hospital design, and especially to consider how improved hospital design can help reduce staff stress and fatigue and increase effectiveness in delivering care, improve patient safety, reduce patient and family stress and improve outcomes and improve overall healthcare quality.

Just as medicine has increasingly moved toward “evidence-based medicine,” where clinical choices are informed by research, healthcare design is increasingly guided by rigorous research linking the physical environment of hospitals to patients and staff outcomes and is moving toward “evidence-based design” (Hamilton, 2003). This report assesses the state of the science that links characteristics of the physical setting to patient and staff outcomes:

- What can research tell us about “good” and “bad” hospital design?
- Is there compelling scientifically credible evidence that design genuinely impacts staff and clinical outcomes?
- Can improved design make hospitals less risky and stressful for patients, their families, and for staff?

In this project, research teams from Texas A&M University and Georgia Tech combed through several thousand scientific articles and identified more than 600 studies—most in top peer-reviewed journals—that establish how hospital design can impact clinical

outcomes. The team found scientific studies that document the impact of a range of design characteristics, such as single-rooms versus multi-bed rooms, reduced noise, improved lighting, better ventilation, better ergonomic designs, supportive workplaces and improved layout that can help reduce errors, reduce stress, improve sleep, reduce pain and drugs, and improve other outcomes. The team discovered that, not only is there a very large body of evidence to guide hospital design, but a very strong one. A growing scientific literature is confirming that the conventional ways that hospitals are designed contributes to stress and danger, or more positively, that this level of risk and stress is unnecessary: improved physical settings can be an important tool in making hospitals safer, more healing, and better places to work.

## **RESEARCH PROCESS**

The research teams searched through scores of databases and in libraries at Texas A&M, Georgia Institute of Technology, University of Michigan, and elsewhere. The team was looking for studies that are:

- Rigorous, in that they use appropriate research methods that allow reasonable comparisons, and discarding of alternative hypotheses. The research studies were assessed on their rigor, quality of research design, sample sizes, and degree of control.
- High impact, in that the outcomes they explore are of importance to healthcare decision-makers, patients, clinicians, and society.

In 1998, Haya Rubin and her colleagues Amanda Owens and Greta Golden found 84 studies produced since 1968 that met similar criteria (Rubin, Owens, & Golden, 1998). Reviewing the research literature six years later, the team estimated that they would find around 125 rigorous studies. We found more than 600.

## **RESULTS**

The research team found rigorous studies that link the physical environment to patient and staff outcomes in four areas:

1. Reduce staff stress and fatigue and increase effectiveness in delivering care
2. Improve patient safety
3. Reduce stress and improve outcomes
4. Improve overall healthcare quality

### **I. Reduce Staff Stress and Fatigue and Increase Effectiveness in Delivering Care**

There is a growing nurse shortage, and this directly threatens patient safety. And the existing hospital-based nursing force is aging. Registered nurses in the United States average more than 43 years old and will average 50 by 2010 and have a turnover rate

averaging 20 percent per year (JCAHO, 2002). The Joint Commission on Accreditation of Healthcare Organizations (JCAHO), in their 2002 report, *Health Care at the Crossroads: Strategies for Addressing the Evolving Nursing Crisis*, noted that a shortage of nurses in America's hospitals is putting patient lives in danger. JCAHO examined 1,609 hospital reports of patient deaths and injuries since 1996 and found that low nursing staff levels were a contributing factor in 24 percent of the cases. The JCAHO report and surveys of nursing found that physical working conditions, along with support and compensation, are key contributors to turnover and burnout. Environmental support for work has become more critical as the typical patient is more seriously ill, patient loads increase, technology changes, and documentation requirements increase.

While reducing staff stress and fatigue through a healing and supportive environment seems like an obvious goal, there are relatively few studies that have dealt with this issue in any detail. More attention has been given to patient outcomes. However, the following sections review several studies that do document (a) effects of healthcare environments on staff health and safety and (b) improving workplaces to increase staff effectiveness, reduce errors, and increase staff satisfaction. Each section also suggests directions for further research.

### ***Improve Staff Health and Safety through Environmental Measures***

Nurses, physicians, and other healthcare employees work under extremely stressful physical conditions. Several studies deal with healthcare employees' risk of contracting infectious diseases from patients due to airborne and surface contamination (Jiang et al., 2003; Kromhout et al., 2000; Kumari et al., 1998; Smedbold et al., 2002). A recent study conducted in the wake of the SARS epidemic in China found that isolating SARS cases in wards with good ventilation could reduce the viral load of the ward and might be the key to preventing outbreaks of SARS among healthcare workers, along with strict personal protection measures in isolation units (Jiang et al., 2003). Another study in Norway found correlations between environmental factors and nasal symptoms of 115 females who worked at 36 geriatric nursing departments. They found significant decrease in nasal inflammation in relation to presence of *Aspergillus fumigatus* in ventilation supply and elevated room temperatures (Smedbold et al., 2002). An evaluation of 17 acute-care or university hospitals in Canada shows that tuberculosis (TB) infection among healthcare workers was associated with ventilation of general or nonisolation patient rooms of less than two air exchanges per hour. The evaluation included all personnel who worked at least two days per week in the respiratory and physiotherapy departments (Menziés, Fanning, Yuan, & Fitzgerald, 2000). This study, like the others, supports the importance of adequate ventilation with good maintenance for ensuring both staff and patient safety in hospitals. Several good studies demonstrating the risk of the sick building syndrome in hospitals also have been compiled within the 2003 *Guidelines for Environmental Infection Control in Healthcare Facilities* by the US Centers for Disease Control and Prevention (CDC) and the Healthcare Infection Control Practices Advisory Committee (HICPAC) (Schulster & Chinn, 2003).

Nursing staff members are also open to risk of injury from medical equipment such as high-intensity surgical-light sources. One study found that a light source used during surgery could potentially cause retinal damage in surgical staff (Fox & Henson, 1996). Much research has examined the effects of noise on patients, but comparatively few studies are available for healthcare staff. There is evidence that staff perceive higher sound levels as stressful (Bayo, Garcia, & Garcia, 1995; Norbeck, 1985). Importantly, noise-induced stress in nurses correlates with reported emotional exhaustion or burnout (Topf & Dillon, 1988). A recent study by Blomkvist et al. (in press, 2004) examined the effects of higher versus lower noise levels on the same group of coronary intensive-care nurses over a period of months. Lower noise levels were linked with a number of positive effects on staff, including reduced perceived work demands, increased workplace social support, improved quality of care for patients, and better speech intelligibility.

Poor ergonomic design of patient beds and nurses' stations leads to back stress, fatigue, and other injuries among nursing staff. In one nursing home study, Garg and Owen evaluated manual tasks deemed stressful by nursing staff and used the information to select patient-transferring devices and to modify toilets and shower rooms. This nursing intervention resulted in a reduction of back injuries of almost 50 percent, from 83 per 200,000 work hours to 47 per 200,000 work hours. Also, in the four months after the intervention, there were no injuries resulting in lost or restricted workdays (Garg & Owen, 1992).

Thus, reducing staff stress by ergonomic interventions, as well as careful consideration of other issues such as air quality, noise, and light, can have significant impact on staff health. In addition, it is also likely to send a message that maintaining health and safety of staff members is an important goal for the organization.

***Increase Staff Effectiveness, Reduce Errors, and Increase Staff Satisfaction by Designing Better Workplaces***

Jobs by nurses, physicians, and others often require a complex choreography of direct patient care, critical communications, charting, filling meds, access to technology and information, and other tasks. Many hospital settings have not been rethought as jobs have changed, and, as a result, the design of hospitals often increases staff stress and reduces their effectiveness in delivering care. While much research in the hospital setting has been aimed at patients, there is a growing and convincing body of evidence suggesting that improved designs can make the jobs of staff much easier.

Nurses spend a lot of time walking. According to one study, almost 28.9 percent of nursing staff time was spent walking (Burgio, Engel, Hawkins, McCorick, & Scheve, 1990). This came second only to patient-care activities, which accounted for 56.9 percent of observed behavior. At least four studies have shown that the type of unit layout (e.g. radial, single corridor, double corridor) influences amount of walking among nursing staff (Shepley, 2002; Shepley & Davies, 2003; Sturdavant, 1960; Trites, Galbraith, Sturdavant, & Leckwart, 1970), and two studies showed that time saved walking was translated into more time spent on patient-care activities and interaction with family

members. Sturdavant (1960) found that fewer trips were made to patient rooms in radial units as nurses were able to better supervise patients visually from the nursing station, though the average time spent with patients was the same in radial as well as single-corridor designs. Shepley and colleagues (2003) found that nursing staff in the radial unit walked significantly less than staff in the rectangular unit (4.7 steps per minute versus 7.9 steps per minute). However, Shepley and her colleagues noted that radial designs might provide less flexibility in managing patient loads. Trites and colleagues (1970) found that a decrease in the percentage of time spent walking by staff in radial units was correlated to an increase in the percentage of time spent in patient-care activities. Also, the majority of the staff surveyed preferred to work in the radial units. Hendrich's research showed that decentralized nurses stations reduced staff walking and increased patient-care time, especially when supplies also were decentralized and placed near the nurses' station (Hendrich, 2003; IOM, 2004). Centralized location of supplies, however, could double staff walking and substantially reduce care time irrespective of whether nurses stations were decentralized (Hendrich, 2003)

Workplace design that reflects a closer alignment of work patterns and the physical setting, such as redesign of a pharmacy layout, has been shown to improve work flow and reduce waiting times, as well as increase patient satisfaction with the service (Pierce, Rogers, Sharp, & Musulin, 1990). Other studies that compared delivery times in decentralized and centralized pharmacy systems found medication delivery times are reduced by more than 50 percent by using decentralized drug-dose distribution systems (Hibbard, Bosso, Sward, & Baum, 1981; Reynolds, Johnson, & Longe, 1978).

Other aspects of the environment, such as lighting levels and auditory or visual distractions, can also affect staff effectiveness while performing critical tasks such as dispensing medical prescriptions. There are relatively few studies that have examined the effect of environmental factors on medication errors, though this is clearly an area of great interest and future research potential. This is discussed in greater detail in the following section.

## **II. Improve Patient Safety**

### ***Hospital-Acquired Infections***

One critically important way that evidence-based design improves safety is by reducing risk from hospital-acquired infections. The research team identified more than 120 studies linking infection to the built environment of the hospital. Transmission of infection to patients occurs through two general routes: airborne and contact. The research literature shows that the design of the physical environment strongly impacts hospital-acquired infection rates by affecting both airborne and contact transmission routes. The literature suggests a clear pattern wherein infection rates are lower when there is very good air quality and patients are in single-bed rather than multi-bed rooms. Also, there is some evidence that providing numerous, easily accessible alcohol-based hand-rub

dispensers or hand washing sinks can increase hand washing compliance and thereby reduce contact contamination.

### ***Reducing Infections Caused by Airborne Pathogens***

Evidence from many studies leaves no doubt that hospital air quality and ventilation play decisive roles in affecting air concentrations of pathogens such as fungal spores (*Aspergillus*) and, in this way, have major effects on infection rates.

***Source of airborne infections:*** Well-conducted research has linked all of the following to air quality and infection rates: type of air filter, direction of airflow and air pressure, air changes per hour in room, humidity, and ventilation system cleaning and maintenance (Lutz, 2003; McDonald et al., 1998). For example, in one study where six patients and one nurse were involved with an outbreak of epidemic methicillin-resistant *Staphylococcus aureus* (EMRSA-15), an environmental source was suspected, and the ventilation grilles in two patient bays were found to be harboring EMRSA-15 (Kumari et al., 1998). The ventilation system, at that time, was working on an intermittent cycle from 4:00 p.m. to 8:00 p.m. Daily shutdown of the system created negative pressure, sucking air in from the ward environment into the ventilation system and contaminating the outlet grilles. The contaminated air blew back into the ward when the ventilation system was started. In another case, the source of infection was the exhaust ducting of the adjacent isolation room ventilation system that allowed the contaminants to enter the unit via a partially open window positioned above a particular bed.

Several studies have identified hospital construction and renovation activities as the sources of airborne infection outbreaks due to dust or particulate generation (Humphreys et al., 1991; Iwen, Davis, Reed, Winfield, & Hinrichs, 1994; Loo et al., 1996; Opal et al., 1986; Oren, Haddad, Finkelstein, & Rowe, 2001). In one study, high spore counts were found within and outside construction sites in a hospital. After control measures were instituted, no further cases of disseminated aspergillosis were identified (Opal et al., 1986). In another study, a nosocomial (hospital-induced) outbreak of invasive pulmonary aspergillosis (IPA) occurred in acute leukemia patients treated in a regular ward with natural ventilation during extensive hospital construction and renovation. The observed infection rate was 50 percent. At this point, some of the patients were moved to a new hematology ward with high-efficiency particulate air (HEPA) filters. During the following three years, none of the patients hospitalized exclusively in the hematology ward developed IPA, although 29% of leukemia patients still housed in the regular ward contracted IPA (Oren et al., 2001).

***Controlling and preventing airborne infection:*** There is convincing evidence that immunocompromised and other high-acuity patient groups have lower incidence of infection when housed in a HEPA-filtered isolation room (Passweg et al., 1998; Sherertz, et al., 1987; Sherertz & Sullivan, 1985). In one study, bone-marrow transplant recipients were found to have a tenfold greater incidence of nosocomial *Aspergillus* infection, compared to other immunocompromised patient populations, when assigned beds outside of a HEPA-filtered environment (Sherertz, et al., 1987). Air contamination is least in

laminar airflow rooms with HEPA filters, and this approach is recommended for operating-room suites and areas with ultraclean room requirements such as those housing immunocompromised patient populations (Alberti et al., 2001; Arlet, Gluckman, Gerber, Perol, & Hirsch, 1989; Dharan & Pittet, 2002; Friberg, Ardnor, & Lundholm, 2003; Hahn et al., 2002; Sherertz, et al., 1987). (Laminar flows are very even, smooth, low velocity airflows that are used in cleanrooms and other settings where high quality ventilation is critical. But laminar flows are relatively expensive and difficult to achieve because furnishings, vents and other features can create turbulence.) HEPA filters are suggested for healthcare facilities by the CDC and HICPAC, but are either required or strongly recommended in all construction and renovation areas (Sehulster & Chinn, 2003).

Effective prevention or control measures during construction and renovation activities include, for example, portable HEPA filters, installing barriers between the patient care and construction areas, negative air pressure in construction/renovation areas relative to patient-care spaces, and sealing patient windows. There is strong evidence of the impact of using HEPA filters for air intakes near construction and renovation sites (Loo et al., 1996; Mahieu, De Dooy, Van Laer, Jansens, & Ieven, 2000; Opal et al., 1986; Oren et al., 2001). A strong study by Humphreys (1991) demonstrates that HEPA filters are not *by themselves* an adequate control measure, and must be employed in conjunction with other measures such as enhanced cleaning, the sealing of windows, and barriers. Cornet et al. (1999) concludes that carefully directed airflow (e.g. laminar airflow) is important, however, we were unable to find and document cost-benefit analysis in the literature to justify the expense versus effectiveness of laminar airflow for patient-care areas near construction and renovation sites

### ***Reducing Infections by Increasing Hand Washing***

Although infection caused by airborne transmission poses a major safety problem, most infections are now acquired in the hospital via the contact pathway (Bauer, Ofner, Just, Just, & Daschner, 1990; IOM, 2004). It is well-established that the hands of healthcare staff are the principal cause of contact transmission from patient to patient (Larson, 1988). The importance of assiduous hand washing by healthcare workers, accordingly, cannot be overemphasized for reducing hospital-acquired infections. In this context, the fact that rates of hand washing by healthcare staff are low represents a very serious patient safety challenge. Several studies of hand washing in high-acuity units with vulnerable patients have found that as few as one in seven staff members wash their hands between patients: compliance rates in the range of 15 percent to 35 percent are typical; rates above 40 percent to 50 percent are the exception (Albert & Condie, 1981; Graham, 1990). Hand washing compliance tends to be consistently lower in units that are understaffed and have a high patient census or bed-occupancy rate (Archibald, Manning, Bell, Banerjee, & Jarvis, 1997).

Education programs to increase hand washing adherence have yielded disappointing or, at best, mixed results. Some investigations have found that education interventions generate no increase at all in hand washing. Even intensive education or training



programs (classes, group feedback, for example) produce only transient increases in hand washing (Conly, Hill, Ross, Lertzman, & Louie, 1989; Dorsey, Cydulka, & Emerman, 1996; Dubbert, Dolce, Richter, Miller, & Chapman, 1990). Given the tremendous morbidity and mortality associated with high rates of hospital-acquired infections, there is an urgent need to identify more effective ways for producing sustained increases in hand washing. Is there evidence suggesting improved hospital design can be effective in elevating hand washing?

***Effect of number and location of sinks/dispensers on hand washing:*** The research team identified six studies that examined whether hand washing is improved by increasing the ratio of the number of sinks or hand-cleaner dispensers to beds and/or by placing sinks or hand-cleaner dispensers in more accessible locations (Cohen, Saiman, Cimiotti, & Larson, 2003; Graham, 1990; Kaplan & McGuckin, 1986; Muto, Sstrom, & Farr, 2000; Pittet et al., 2000; Vernon, Trick, Welbel, Peterson, & Weinstein, 2003). These studies, on balance, offer support, though limited, for the notion that providing numerous, conveniently located alcohol-rub dispensers or washing sinks can increase compliance. In particular, the evidence suggests that installing alcohol-based hand-cleaner dispensers at bedside usually improves adherence. As an example, a study by Pittet et al. (2000) found that a combination of bedside antiseptic hand-rub dispensers and posters to remind staff to clean their hands substantially increased compliance. Cohen et al. (2003) likewise reported improved adherence in association with the installation of numerous alcohol-gel dispensers. By contrast, Muto et al. (2000) reported that placing alcohol-gel dispensers next to the doors of patient rooms did not increase adherence. Two other investigations focusing on sinks (water/soap) identified a positive relationship between observed frequency of hand washing and a higher ratio of sinks to beds (Kaplan & McGuckin, 1986; Vernon et al., 2003). Providing automated water/soap sinks, however, appears not to increase hand washing rates compared to traditional non-automated sinks (Larson, et al., 1991; Larson, Bryan, Adler, & Blane, 1997).

Further, three studies offer convincing and important evidence that providing single-patient rooms with a conveniently located sink in each room reduces nosocomial infection rates in intensive care units, such as neonatal intensive care (NICU) or burn units, compared to when the same staff and comparable patients are in multibed open units with few sinks (Goldmann, Durbin, & Freeman, 1981; McManus, A. T., Mason, McManus, & Pruitt, 1994; McManus, A. T., McManus, Mason, Aitcheson, & Pruitt, 1985; Mulin et al., 1997). Although hand washing frequency was not measured in these studies, the investigators posited that increased hand washing was an important factor in reducing infections in the units with single rooms and more sinks. A comparison of an ICU converted from an open unit with few sinks to single rooms with one sink per room found a nonsignificant tendency for hand washing to increase (from 16 percent to 30 percent) but no decline in infection incidence (Preston, Larson, & Stamm, 1981). These results are perhaps explainable by the fact that several sinks in the single-bed unit were placed in comparatively inaccessible or inconvenient locations, such as behind doors or away from staff work paths.

Despite the encouraging overall pattern of findings in these studies, it is not clear how much of the effectiveness in terms of increased hand washing or reduced infection rates can be attributed to the installation of more numerous and/or accessible sinks and alcohol-gel dispensers. Future research should include controlled experiments that systematically vary the number and location of hand-cleaning stations or dispensers. There is also a conspicuous need for studies that define *accessible* locations for hand-cleaning stations in an evidence-based manner—that is, on the basis of empirical analysis of staff movement paths, visual fields, interactions with patients and families, and work processes. In this regard, the neglect of *human factors* knowledge and research methods is a major weakness of the hand washing research and, more generally, of the infection control literature. Research teams should include a human factors specialist and often an environmental psychologist. The urgent need to increase hand washing frequency underscores the high priority that should be accorded this research direction.

### ***Reducing Infections with Single-Bed Rooms***

The research team identified at least 16 studies relevant to the question of whether nosocomial infection rates differ between single-bed and multi-bed rooms. The findings collectively provide a strong pattern of evidence indicating that infection rates are usually lower in single-bed rooms. Different mechanisms or factors have been identified or implicated as contributing to lower infection incidence in single rooms. One clear set of advantages relates to reducing airborne transmission through air quality and ventilation measures such as HEPA filters, negative room pressure to prevent a patient with an aerial-spread infection from infecting others, or maintaining positive pressure to protect an immunocompromised patient from airborne pathogens in nearby rooms. A strong study by Passweg et al. (1998) found that the combination of room isolation and HEPA filtration reduced infection and mortality in bone marrow transplant patients and significantly increased their one-year survival rates. Research studying burn patients also has shown that single rooms and good air quality substantially reduce infection incidence and reduce mortality (McManus, A. T. et al., 1994; McManus, A.T., Mason, McManus, & Pruitt, 1992; Shirani et al., 1986; Thompson, Meredith, & Molnar, 2002). Studies of cross-infection for contagious airborne diseases (influenza, measles, TB, for example) have found, as would be expected, that placing patients in single rooms is safer than housing them in multibed spaces (Gardner, Court, Brocklebank, Downham, & Weightman, 1973; McKendrick & Emond, 1976).

Severe Acute Respiratory Syndrome (SARS) outbreaks in Asia and Canada dramatically highlighted the shortcomings of multibed rooms for controlling or preventing infections both for patients and healthcare workers. SARS is transmitted by droplets that can be airborne over limited areas. Approximately 75 percent of SARS cases in Toronto resulted from exposure in hospital settings (Farquharson & Baguley, 2003). The pervasiveness in Canadian and Asian hospitals of multibed spaces in emergency departments and ICUs, together with the scarcity of isolation rooms with negative pressure, severely hindered treatment and control measures. Toronto hospitals were forced to create additional negative-pressure isolation rooms by quickly constructing wall barriers to replace bed curtains and making airflow and pressure adaptations (Farquharson & Baguley, 2003).

In addition to clear advantages in reducing airborne transmission, several studies show that single-bed rooms also lessen risk of infections acquired by contact. As background for understanding how single rooms can lessen contact spread, it should first be mentioned that many environmental surfaces and features become contaminated near infected patients. Examples of surfaces found to be contaminated frequently via contact with patients and staff include: overbed tables, bed privacy curtains, computer keyboards, infusion pump buttons, door handles, bedside rails, blood pressure cuffs, chairs and other furniture, and countertops (Aygün et al., 2002; Boyce, Potter-Bynoe, Chenevert, & King, 1997; Bures, Fishbain, Uyehara, Parker, & Berg, 2000; Devine, Cooke, & Wright, 2001; Neely & Maley, 2001; Noskin, Bednarz, Suriano, Reiner, & Peterson, 2000; Palmer, 1999; Roberts, Findlay, & Lang, 2001; Rountree, Beard, Loewenthal, May, & Renwick, 1967; Sanderson & Weissler, 1992; Williams, Singh, & Romberg, 2003). These and other contaminated surfaces and features act as pathogen reservoirs that increase cross-infection risk. Boyce et al. (1997) found that in the rooms of patients infected with methicillin-resistant *Staphylococcus aureus* (MRSA), 27 percent of all environmental surfaces sampled were contaminated with MRSA.

Compared to single-bed rooms, multi-bed rooms are far more difficult to decontaminate thoroughly after a patient is discharged, and therefore worsen the problem of multiple surfaces acting as pathogen reservoirs. Because different staff members who enter a room can touch the same contaminated surfaces, the risk of a nurse unknowingly becoming contaminated should be greater in multioccupancy rooms. Circumstantial support for this point is provided by research on contamination of nurses in units having patients infected by MRSA. Boyce et al. (1997) found that 42 percent of nurses who had no direct contact with an MRSA patient but had touched contaminated surfaces contaminated their gloves with MRSA.

In a study of MRSA infections in NICUs, Jernigan et al. (1996) reported that risk was lowered by isolation in single-bed rooms; high risk was associated with spatial proximity to an infected patient and shared exposure to caregivers. Ben-Abraham et al. (2002) found that nosocomial infection frequency was much lower in a single-bed pediatric intensive care unit than a unit with multi-bed rooms. The investigators tentatively concluded that single-bed rooms helped to limit person-to-person spread of pathogens between pediatric patients. Having a roommate has been identified as a risk factor for nosocomial diarrhea and gastroenteritis (Chang, V. T. & Nelson, 2000; Pegues & Woernle, 1993).

To summarize briefly, there is a convincing pattern of evidence across many studies indicating that single-bed rooms lower nosocomial infection rates. Singles appear to limit person-to-person and person-surface-person spread of infection in part because they are far easier to decontaminate thoroughly than multibed rooms after patients are discharged. Also, single rooms with a conveniently located sink or alcohol-gel dispenser in each room may heighten hand washing compliance compared to multibed rooms with few sinks. Finally, single rooms are clearly superior to multi-bed rooms with respect to reducing airborne transmission of pathogens.

### ***Reducing Medication Errors***

The research team identified three rigorous studies that link environmental factors, such as lighting, distractions, and interruptions, with errors in prescribing or dispensing medications (Booker & Roseman, 1995; Buchanan, Barker, Gibson, Jiang, & Pearson, 1991; Flynn et al., 1999). Although there are relatively few studies in this area, the findings suggest a promising research direction that merits further exploration, considering the vital importance of preventing medical errors in hospitals. Results from a large-scale study of the effects of different illumination levels on pharmacists' prescription-dispensing error rates strongly suggested that such errors are reduced when work-surface light levels are relatively high (Buchanan et al., 1991). In this study, three different illumination levels were evaluated (450 lux, 1,100 lux, 1,500 lux). Medication-dispensing error rates were significantly lower (2.6 percent) at an illumination level of 1,500 lux (highest level), compared to an error rate of 3.8 percent at 450 lux. Two investigations of medication dispensing errors by hospital pharmacists found that error rates increased sharply for prescriptions when an interruption or distraction occurred, such as a telephone call (Flynn et al., 1999; Kistner, Keith, Sergeant, & Hokanson, 1994). Thus, lighting levels, frequent interruptions or distractions during work, and inadequate private space for performing work can be expected to worsen medication errors. The process of improving systems and processes to reduce medication errors in hospitals should include an assessment of the environment in which staff members perform their activities. Additional research is required to confirm the findings from studies as well as to identify ways to design better working environments that may reduce or prevent the likelihood of such errors occurring.

There is mounting evidence that the transfer of patients between rooms or different units is a source of medication errors (Cook, Render, & Woods, 2000). Reasons why errors plague room transfers include delays, communication discontinuities among staff, loss of information, and changes in computers or systems. The solution implied is to create an acuity-adaptable care process and patient rooms that substantially reduce transfers. When Methodist Hospital in Indianapolis, Indiana, changed from two-bed rooms in coronary intensive care to acuity-adaptable single-bed rooms, transfers declined 90 percent and medication errors were lowered by 67 percent (Hendrich, Fay, & Sorrells, 2002; 2004). Reducing transfers also saves much staff time, shortens patient stays, and reduces costs (IOM, 2004). Further studies and demonstration projects are needed to ascertain the safety advantages of acuity-adaptable rooms for other types of units and patient categories.

### ***Reduce Patient Falls***

There is a very large literature that looks at the causes and risk factors involved in patient falls in hospitals. This is an area of great importance because patients who fall incur physical injuries, psychological effects, and have greater lengths of stay in the hospital (Brandis, 1999). It is estimated that the total cost of fall injuries for older people was

around \$20.2 billion per year in the United States in 1994, and is projected to reach \$32.4 billion (in 1994 U.S. dollars) in 2020 (Chang, 2004). While the role of the environment in causing or preventing patient falls is widely accepted, there is not yet evidence conclusively tying environmental interventions with reduced falls. Available studies usually examine the location of fall incidents retrospectively or discuss environmental-modification programs such as improving lighting, securing carpeting, and so on. However, a meta-analysis and systematic review of randomized controlled trials of fall-prevention interventions found that there was no clear evidence for the independent effectiveness of environmental-modification programs (Chang, 2004).

Nonetheless, several studies have shown that most patient falls occur in the bedroom, followed by the bathroom, and that comprehensive fall-prevention programs can have an effect. Brandis (1999) reported transfers to and from bed as the cause of 42.2 percent of inpatient falls. Design faults identified in the bathroom and bedroom areas included slippery floors, inappropriate door openings, poor placement of rails and accessories, and incorrect toilet and furniture heights. After the fall-prevention program (which included identifying high-risk patients, management strategies, environmental and equipment modification, and standardization) was implemented, there was an overall decrease of 17.3 percent in falls. Thus, fall-prevention strategies that have included environmental modification have worked in the past. However, it is not clear how much of the effectiveness of such strategies can be attributed to environmental factors alone.

An innovative and promising environmental strategy for reducing falls has its origins in evidence suggesting that many falls occur when patients attempt to get out of bed unassisted or unobserved (Uden, 1985; Vassallo, Azeem, Pirwani, Sharma, & Allen, 2000). It should be mentioned that considerable evidence has shown that bedrails are ineffective for reducing the incidence of falls and may increase the severity of fall injuries from beds (Capezuti, Maislin, Strumpf, & Evans, 2002; Hanger, Ball, & Wood, 1999; van Leeuwen, Bennett, West, Wiles, & Grasso, 2001). To increase observation and improve assistance for patients and thereby reduce falls, Methodist Hospital in Indianapolis, Indiana, changed from a coronary critical care unit with centralized nurses stations and two-bed rooms to one having decentralized nurses stations and large single-bed rooms designed to support family presence (Hendrich, et al., 2002). Comparison of data from two years prior and three years after the new unit design showed that falls were cut by 2/3—from six-per-thousand patients to two-per-thousand. Given that falls are a critical safety problem, additional research is needed to understand the effectiveness of this approach for designing patient-care units.

### ***Improve Patient Confidentiality and Privacy***

Confidentiality has emerged as a priority issue in light of research showing that physicians and nurses very frequently breach patient confidentiality and privacy by talking in spaces where they are overhead by other patients or persons (Ubel, Zell, & Miller, 1995). The seriousness of the problem is underscored, for example, by a study of an emergency department at a university hospital that showed that 100 percent of physicians and other clinical personnel committed confidentiality and privacy breaches

(Mlinek & Pierce, 1997). HIPAA, the Health Insurance Portability and Accountability Act of 1998, has further elevated the importance of providing reasonable safeguards to protect the confidentiality of staff conversations with and about patients.

Although the importance of the built environment for patient confidentiality may seem self-evident, only a few studies have directly examined the role of unit design or architecture. A study by Barlas et al. (2001) compared auditory and visual privacy for emergency department patients assigned to either multibed spaces with curtain partitions or rooms with solid walls. Those with curtains reported having far less auditory and visual privacy than patients with walls. An important finding was that 5 percent of the patients in curtained spaces reported they withheld portions of their medical history and refused parts of their physical examination because of lack of privacy (Barlas et al., 2001). None of the patients in rooms with walls reported withholding information. The fact that some emergency department patients with curtains withheld information suggests that lack of privacy can reduce patient safety. Additional convincing evidence of the importance of the emergency department physical environment comes from a study that documented frequent breaches of auditory and visual privacy and confidentiality in areas with curtains compared to rooms with solid walls (Mlinek & Pierce, 1997). A recent questionnaire study of staff in four West Coast hospitals found that nurses overwhelmingly judged single rooms to be superior to double rooms for examining a patient (85 percent) and for collecting a patient's history (82 percent) (Chaudhury, Mahmood, & Valente, 2003).

There is a clear need for additional studies that examine privacy and confidentiality breaches associated with the physical environment in single versus double rooms, multibed spaces in ICUs and other types of units, and in spaces such as waiting rooms and nurses' stations. As well, priority should be accorded to generating more research that investigates how the quality of communication and information *from* patients *to* physicians and nurses is affected by the unit architecture.

Analysis of patient satisfaction data made available by Press Ganey (2003) for this report leaves no doubt that patients in single-bed rooms, compared to those with a roommate, are consistently much more satisfied with "concern for your privacy." The satisfaction data were obtained from 2,122,439 patients who received care during 2003 in 1,462 healthcare facilities (Press Ganey, 2003). Fifty-six percent of the patients were in single rooms; 44 percent had a roommate. Greater satisfaction with privacy in single rooms was evident across all major patient categories and types of unit and across different age and gender groups. Satisfaction with privacy was 4.5 percent higher on average nationally in single rooms than doubles—a substantial difference considering that it can be difficult for hospitals to increase satisfaction scores by even 1 percent to 2 percent.

### III. Reduce Stress and Improve Outcomes

#### *Reduce Noise*

The research team identified more than 130 references focusing on noise in hospitals in the research literature. Studies tend to fall into three broad categories: those that measure noise levels in hospital spaces but do not assess outcomes, studies of the effectiveness of environmental and/or organizational interventions in reducing noise, and investigations of the effects of noise on outcomes.

World Health Organization guideline values for continuous background noise in hospital patient rooms are 35 dB, with nighttime peaks in wards not to exceed 40 dB (Berglund, Lindvall, & Schwela, 1999). These guidelines notwithstanding, many studies have shown that hospital background noise levels fall in far higher ranges. Background noise levels typically are 45 dB to 68 dB, with peaks frequently exceeding 85 dB to 90 dB (Aaron et al., 1996; Allaouchiche, Duflo, Debon, Bergeret, & Chassard, 2002; Blomkvist et al., in press, 2004; Falk & Woods, 1973; Hilton, 1985; McLaughlin, McLaughlin, Elliott, & Campalani, 1996; Robertson, Cooper-Peel, & Vos, 1998). In judging these noise levels, it is worth noting that the decibel scale is logarithmic; each 10 dB increase represents approximately a doubling in the perceived sound level. A 60 dB sound, accordingly, is perceived as roughly four times as loud as a 40 dB sound. Medical equipment and staff voices often produce 70 dB to 75 dB levels measured at the patient's head, which approach the noise level in a busy restaurant (Blomkvist et al., in press, 2004). Noises from alarms and certain equipment exceed 90 dB (for example, portable X-ray machine), which is comparable to walking next to a busy highway when a motorcycle or large truck passes. A study in a NICU measured peak levels once per minute and found that 31 percent of peaks exceed 90 dB (Robertson et al., 1998). Noise peaks in hospitals can be extraordinarily loud. A recent study recorded 113 dB during shift changes at a large hospital (Cmiel, Karr, Gasser, Oliphant, & Neveau, 2004). Operating room noises from drills, saws, and other equipment are in the range of 100 dB to 110 dB, presenting a significant risk for noise-induced hearing loss (Hodge & Thompson, 1990; Love, 2003; Nott & West, 2003).

The research reviewed suggests that hospitals are excessively noisy for two general reasons (Ulrich, Lawson, & Martinez, 2003). First, noise sources are numerous, often unnecessarily so, and many are loud. Well-documented examples include paging systems, alarms, bedrails moved up/down, telephones, staff voices, ice machines, pneumatic tubes, trolleys, and noises generated by roommates. Second, environmental surfaces—floors, walls, ceilings—usually are hard and sound-reflecting, not sound-absorbing, creating poor acoustic conditions. Sound-reflecting surfaces cause noise to propagate considerable distances, traveling down corridors and into patient rooms, and adversely affecting patients and staff over larger areas. Sound-reflecting surfaces typical of hospitals cause sounds to echo, overlap, and linger or have long reverberation times (Blomkvist et al., in press, 2004; Ulrich et al., 2003).

Environmental interventions that have proven especially effective for reducing noise and improving acoustics in hospital settings include: installing high-performance sound-absorbing ceiling tiles, eliminating or reducing noise sources (for example, adopting a noiseless paging system), and providing single-bed rather than multibed rooms. In general, studies of the effectiveness of noise-reduction measures suggest that environmental or design interventions are more successful than organizational interventions such as staff education or establishing quiet hours (Gast & Baker, 1989; Moore, 1998; Walder, Francioli, Meyer, Lancon, & Romand, 2000).

A clear-cut finding in the literature is that noise levels are much lower in single-bed than multi-bed rooms. Studies of multi-bed rooms in acute care and intensive care units have shown that most noises stem from the presence of another patient (staff talking, staff caring for other patients, equipment, visitors, patient sounds such as coughing, crying out, rattling bed rails) (Baker, 1984; Southwell & Wistow, 1995; Yinnon, Ilan, Tadmor, Altarescu, & Hershko, 1992). A study of multi-bed bays in a children's hospital concluded that noise levels were so high that consideration should be given to abolishing open bay rooms (Couper et al., 1994). Further, patient satisfaction data provided for this report by Press Ganey (2003) unequivocally show that patients in single-bed rooms, compared to those with a roommate, are vastly more satisfied with the "noise levels in and around your room." The satisfaction data, as noted in an earlier section, were obtained from 2,122,439 patients who received care during 2003 in 1,462 healthcare facilities (Press Ganey, 2003). Far higher satisfaction with noise levels in single rooms was evident across all major patient categories and types of unit and across different age and gender groups. Satisfaction with noise level was 11.2 percent higher on average nationally in single rooms than doubles—a huge difference. As was noted above, it can be difficult for hospitals to achieve even 1 percent to 2 percent increases in patient satisfaction scores. The combination of findings from noise-level measurement studies and patient-satisfaction surveys highlight the great advantage of providing single rooms, compared to two-bed rooms, with respect to reducing noise.

A considerable body of research has documented negative effects of noise on patient outcomes. Several studies have focused on infants in NICUs, finding that higher noise levels, for example, decrease oxygen saturation (increasing need for oxygen support therapy), elevate blood pressure, increase heart and respiration rate, and worsen sleep (Johnson, 2001; Slevin, Farrington, Duffy, Daly, & Murphy, 2000; Zahr & de Traversay, 1995). Much research on adults and children has unequivocally shown, as might be expected, that noise is a major cause of awakenings and sleep loss (Blomkvist et al., in press, 2004; Gabor et al., 2003; Meyer et al., 1994; Parthasarathy & Tobin, 2004; Schnelle, Ouslander, Simmons, Alessi, & Gravel, 1993; Topf, 1985; Topf & Davis, 1993; Topf & Thompson, 2001; Yinnon et al., 1992). In multi-bed rooms, noises stemming from the presence of other patients often are the major cause of sleep loss. Berg (2001) found, by monitoring brain activity, that even relatively low decibel levels—38 dB to 40 dB—when coupled with longer reverberation times (sound-reflecting ceiling) significantly fragmented and worsened sleep of volunteers in patient rooms. Berg's (2001) findings have disturbing implications because most hospitals have nighttime sound peaks exceeding those of the patient rooms in his study.



Apart from worsening sleep, there is strong evidence that noise increases stress in adult patients, for example, heightening blood pressure and heart rate (Baker, 1992; Morrison, Haas, Shaffner, Garrett, & Fackler, 2003; Novaes, Aronovich, Ferraz, & Knobel, 1997; Topf & Thompson, 2001). A recent study by Blomkvist et al. (in press, 2004) examined the effects of poor versus good sound levels and acoustics on coronary intensive-care patients by periodically changing the ceiling tiles from sound-reflecting to sound-absorbing tiles. When the sound-absorbing ceiling tiles were in place, patients slept better, were less stressed (lower sympathetic arousal), and reported that nurses gave them better care. There were also indications in this study that the incidence of re-hospitalization was lower if patients had experienced the sound-absorbing rather than sound-reflecting ceiling during their hospital stay (Hagerman et al., in press, 2004). More studies are needed such as that by Blomkvist et al. (in press, 2004), which use experimental research designs and systematically vary noise conditions. Future research should also investigate the effects of noise on re-hospitalization rates and other outcomes. In sum, the main message from the research review is clear: new hospitals should be much quieter, and effective design strategies for quieting hospitals are available.

### *Improve Sleep*

The above section reviewed many studies showing that noise levels are high in hospitals and that noise is a major cause of poorer sleep in patients. Interventions that reduce hospital noise have been found to improve sleep and reduce patient stress. As noted earlier, environmental interventions found to be most effective for reducing noise in hospital settings include: providing single-bed rather than multi-bed rooms, installing high-performance, sound-absorbing ceiling tiles, using sound-absorbing flooring where possible, and eliminating or reducing noise sources (for example, use noiseless paging, locate alarms outside patient rooms).

### *Reduce Spatial Disorientation*

Wayfinding problems in hospitals are costly and stressful and have particular impacts on outpatients and visitors, who are often unfamiliar with the hospital and are otherwise stressed and disoriented. In a study conducted at a major regional 604-bed tertiary-care hospital, the annual cost of the wayfinding system was calculated to be more than \$220,000 per year in the main hospital or \$448 per bed per year in 1990. Much of this cost was the hidden costs of direction giving by people other than information staff, which occupied more than 4,500 staff hours, the equivalent of more than two full-time positions (Zimring, 1990). While almost all hospitals strongly feel the problems associated with a complicated building and poor wayfinding system, it is usually difficult to tackle this problem with a piecemeal approach. A wayfinding system, as the name implies, is not just about better signage or colored lines on floors. Rather, hospitals are seeking to provide integrated systems that include coordinated elements such as visible and easy-to-understand signs and numbers, clear and consistent verbal directions, consistent and clear paper, mail-out and electronic information and a legible physical

setting (Carpman, 1993). A wayfinding system includes four main components that work at different levels: administrative and procedural levels, external building cues, local information and global structure

***Administrative and procedural information:*** Mail-out maps, electronic information available on the Web and at kiosks and verbal directions are organizational strategies aimed at providing key information to patients to prepare them for their hospital visit. This is not dealt with in this review.

***External building cues:*** Signs and cues that lead to the hospital, especially the parking lot, need to be considered carefully, as they are the first point of contact of the patient with the hospital (Carpman, Grant, & Simmons, 1985). For example, Carpman, Grant and Simmons conducted a video simulation study to assess the relative role of signs and seeing a destination. The hospital wanted to direct most traffic to a parking structure rather than a drop-off lane. When the researchers showed prospective visitors a simulated video showing a design alternative that allow arriving drivers to see the main pavilion with the drop-off lane, 37 percent of the respondents said that they would turn into the drop circle when they could see the entry to the garage, ignoring the signs. As a consequence, the hospital chose to redesign the entry.

***Local information:*** Once patients find their way to the building from the parking lot, they are faced with the prospect of identifying the destination. Informational handouts, information desks, you-are here maps, directories, and signage along the way are critical wayfinding aids (Carpman, Grant, & Simmons, 1983-84; Levine, Marchon, & Hanley, 1984; Nelson-Shulman, 1983-84; Wright, Hull, & Lickorish, 1993). In an experimental study, researchers found that patients who had the benefit of an information system (welcome sign, hospital information booklet, patient letter, orientation aids) upon reaching the admitting area were more self-reliant and made fewer demands on staff. In contrast, uninformed patients rated the hospital less favorably and were found to have elevated heart rates (Nelson-Shulman, 1983-84).

Information provided in you-are-here maps can be useful. However, you-are-here maps should be oriented so that the top signifies the direction of movement for ease of use. When the maps were aligned in directions other than the forward position, people not only took much longer to find their destination, but were significantly less accurate (Levine et al., 1984). Another study found that people who used signs found their destination faster than those who only used maps (Butler, Acquino, Hissong, & Scott, 1993). However, people who were given a combination of handheld maps and wall signs reached their destination more often than those who just used wall signs (Wright et al., 1993).

It is critical to design signage systems with logical room numbering and comprehensible nomenclature for departments (Carpman & Grant, 1993; Carpman, Grant, & Simmons, 1984). For example, inpatients, outpatients, and visitors to a hospital preferred simple terms such as *walkway* or *general hospital* over more complex or less-familiar terms such as *overhead link*, *medical pavilion* or *health-sciences complex*.

Contrary to the belief that fewer signs in hospital hallways means less clutter and hence less confusion, an experimental study in a hospital found that patients who had access to more number of signs along the way to the destination were faster, less hesitant, asked for directions a fewer number of times, and reported lower levels of stress (Carpman et al., 1984). Based on this study, the authors suggest that directional signs should be placed at or before every major intersection, at major destinations, and where a single environmental cue or a series of such cues (e.g. change in flooring material) convey the message that the individual is moving from one area into another. If there are no key decision points along a route, signs should be placed approximately every 150 feet to 250 feet.

**Global structure:** In addition to local properties of the spaces that people move through, there are specific characteristics of the overall structure of the system of rooms and corridors that impact the paths people take (Haq & Zimring, 2003; Peponis, Zimring, & Choi, 1990). People tend to move toward spaces and through corridors that are more accessible from a greater number of spaces. Based on observations of search patterns of study participants in a hospital and use of objective measures that quantify spatial characteristics, researchers found that participants tended to move along more “integrated” routes—routes that are, on average, more accessible because they are fewer turns from all other routes in the hospital. This research suggests that it may be important to identify such integrated routes in the plan while placing important facilities and key points such as the entrance (Peponis et al., 1990).

The research supports the value of a systems approach to wayfinding. Wayfinding continues to be a pervasive problem in hospitals because it is not sufficient to consider one or two components separately. Well-designed signs are likely to be quite ineffective in a building that is highly complicated and does not provide simple cues that enable natural movement. While there are more than 17 studies that look at wayfinding in hospitals and other buildings (Brown, Wright, & Brown, 1997; Carpman & Grant, 1993; Carpman et al., 1983-84, 1984; Carpman et al., 1985; Christensen, 1979; Grover, 1971; Haq & Zimring, 2003; Levine et al., 1984; Moeser, 1988; Nelson-Shulman, 1983-84; Ortega-Andeane & Urbina-Soria, 1988; Passini, Rainville, Marchand, & Joannette, 1995; Peponis et al., 1990; Schneider, L. F. & Taylor, 1999; Weisman, 1981; Wright et al., 1993; Zimring & Timpler, 1983-84), it is quite difficult to isolate single influences of design on wayfinding performance or of wayfinding on outpatient or visitor stress. The problem is exacerbated by the fact that most hospitals have existing complex buildings upon which they try to superimpose a signage system to make things work. This strategy is ineffective in most cases.

There are some very good studies that deal with designing better signage, optimal spacing and location of signage, types of information that are most effective in way finding, and so on. Similarly, other studies at the global scale have looked at the properties of the building layout that facilitate or impede movement. It is essential that these different pieces of information come together while designing new hospitals where there is opportunity to develop an effective wayfinding system at multiple levels. Additional

studies are needed to ascertain the magnitude of stress that wayfinding problems have on outpatients and family.

### ***Reduce Depression***

Several studies strongly support that bright light—both natural and artificial—can improve health outcomes such as depression, agitation, sleep, circadian rest-activity rhythms, as well as length of stay in demented patients and persons with seasonal affective disorders (SAD). At least eleven strong studies suggest that bright light is effective in reducing depression among patients with bipolar disorder or SAD. Further, seven studies show that exposure to morning light is more effective than exposure to evening light in reducing depression (Beauchemin & Hays, 1996; Benedetti, F., Colombo, C., Barbini, B., Campori, E., & Smeraldi, E., 2001; Lewy et al., 1998; Lovell, Ancoli-Israel, & Gevirtz, 1995; Terman, Terman, Lo, & Cooper, 2001; Van Someren, Kessler, Mirmiran, & Swaab, 1997; Wallace-Guy et al., 2002). An experimental study that compared the effect of morning and evening light on patients with winter depression found that morning light was *twice* as effective as evening light in treating SAD (Lewy et al., 1998). Exposure to bright morning light has been shown to reduce agitation among elderly patients with dementia. When elderly patients with dementia were exposed to 2,500 lux for two hours in the morning for two ten-day periods, their agitation reduced. Patients were significantly more agitated on non-treatment days (Lovell et al., 1995).

There is also strong evidence that exposure to bright light improves sleep and circadian rhythms. When the daytime environmental illumination level was increased in different living spaces of a dementia unit, it was found that, during increased illumination periods, the stability of the rest-activity rhythm increased in patients with intact vision, but not in visually impaired patients (Van Someren et al., 1997).

It has also been shown that patients in brightly lit rooms have a shorter length of stay compared to patients in dull rooms. Beauchemin and Hays (1996) found that patients hospitalized for severe depression reduced their stays by an average of 3.67 days if assigned to a sunny rather than a dull room overlooking spaces in shadow.

A recent randomized prospective study assessed whether the amount of sunlight in a hospital room modifies a patient's psychosocial health, quantity of analgesic medication used, and pain medication cost (Walch et al., 2004, in press). Patients undergoing elective cervical and lumbar spinal surgeries were admitted to the bright or the dim side of the inpatient surgical ward postoperatively. The outcomes measured included the standard morphine equivalent of all opioid medication used postoperatively by patients and their subsequent pharmacy cost. This strong study found that patients exposed to an increased intensity of sunlight experienced less perceived stress, less pain, took 22 percent less analgesic medication per hour and had 20 percent less pain medication costs.

Using light as an intervention to reduce depression in clinically depressed as well as non-depressed patients is a relatively inexpensive intervention that has been shown to yield consistently positive results. While many studies deal with the effects of artificial light,

natural daylight in patient rooms has also been found effective in reducing depression, reducing length of stay, and reducing intake of pain medication. Thus, an important consideration while designing hospital layouts may be to optimize exposure to morning light in patient rooms by using an east-facing orientation. This research further implies the possibility that depression might be worsened by architectural designs that block or sharply reduce natural daylight in patient rooms. A hypothetical example would be a hospital having patient-room windows looking out into a roofed atrium with few skylights and little natural daylight. In this example, deprivation of natural daylight could be extreme if patient windows were tinted to prevent users of the atrium from looking into patient rooms and violating privacy.

### ***Provide Nature and Positive Distraction***

Positive distractions refer to a small set of environmental features or conditions that have been found by research to effectively reduce stress. Distractions can include certain types of music, companion animals such as dogs or cats, laughter or comedy, certain art, and especially nature (Ulrich, 1991). The focus here is on the last, nature. (There is an additional large research literature on music, but this is not covered in this review.)

As background relevant to assessing the credibility of nature findings in healthcare environments, it should be mentioned that many studies of populations other than hospital patients have produced strong evidence that even fairly brief encounters with real or simulated nature settings can elicit significant recovery from stress within three minutes to five minutes at most (Parsons & Hartig, 2000; Ulrich, 1999). Investigators have consistently reported that stress-reducing or restorative benefits of simply viewing nature are manifested as a constellation of positive emotional and physiological changes. Stressful or negative emotions such as fear or anger diminish while levels of pleasant feelings increase. Laboratory and clinical studies have shown that viewing nature produces stress recovery quickly evident in physiological changes, for instance, in blood pressure and heart activity (Ulrich, 1991). By comparison, considerable research has demonstrated that looking at built scenes lacking nature (rooms, buildings, parking lots) is significantly less effective in fostering restoration and may worsen stress.

Questionnaire studies have found that bedridden patients assign especially high preference to having a hospital window view of nature (Verderber, 1986). Mounting research is providing convincing evidence that visual exposure to nature improves outcomes such as stress and pain. For example, a study in a Swedish hospital found that heart-surgery patients in ICUs who were assigned a picture with a landscape scene with trees and water reported less anxiety/stress and needed fewer strong doses of pain drugs than a control group assigned no pictures (Ulrich, 1991). Another group of patients assigned an abstract picture, however, had worsened outcomes compared to the control group. Ulrich (1984) found that patients recovering from abdominal surgery recovered faster, had better emotional well-being, and required fewer strong pain medications if they had bedside windows with a nature view (looking out onto trees) than if their windows looked out onto a brick wall.

Recently, strong studies using experimental designs have produced additional convincing evidence that viewing nature reduces patient pain as well as stress. These investigations also support the interpretation that nature serves as a positive distraction (Ulrich, 1991) that reduces stress and diverts patients from focusing on their pain or distress. A randomized prospective investigation found that adult patients undergoing a painful bronchoscopy procedure reported less pain if they were assigned to look at a ceiling-mounted nature scene rather than a control condition consisting of a blank ceiling (Diette, Lechtzin, Haponik, Devrotes, & Rubin, 2003). Another controlled experiment that used volunteers in a hospital assessed the effect on pain of viewing a soundless nature videotape in contrast to a static blank screen (Tse, Ng, Chung, & Wong, 2002). Subjects who watched the nature scenes evidenced a higher threshold for detecting pain and had substantially greater pain tolerance. Two studies of female cancer patients have shown that taking a virtual reality nature walk while in bed or a hospital room (through a forest with bird sounds) reduced anxiety and symptomatic distress (Schneider, S. M., Prince-Paul, Allen, Silverman, & Talaba, 2004). Research on patients suffering intense pain because of severe burns found that exposing patients to a videotape of scenic nature (forest, flowers, ocean, waterfalls) during burn dressing changes significantly reduced both anxiety and pain intensity (Miller, Hickman, & Lemasters, 1992).

The possibility that nature can improve outcomes even in patients with late-stage dementia, including Alzheimer's disease, has received some support from a quasi-experimental study that found reduced levels of agitated aggressive behavior associated with a shower bath when recorded nature sounds (birds, babbling brook) and color pictures were present (Whall et al., 1997). A well-controlled study of blood donors in a waiting room found that blood pressure and pulse were lower on days when a wall-mounted television displayed a nature videotape, compared to days with continuous daytime television programs (Ulrich, Simons, & Miles, 2003). More research is needed to identify conditions under which television can either be a stress-reducing positive distraction or a stressor in hospitals.

***Gardens in healthcare environments:*** Hospital gardens not only provide restorative or calming nature views, but can also reduce stress and improve outcomes through other mechanisms, for instance, fostering access to social support and providing opportunities for positive escape and sense of control with respect to stressful clinical settings (Cooper Marcus & Barnes, 1995; Ulrich, 1999). Based on postoccupancy evaluations of four hospital gardens in California, Cooper-Marcus and Barnes (1995) concluded that many nurses and other healthcare workers used the gardens for achieving pleasant escape and recuperation from stress. Other postoccupancy studies indicate that patients and family who use hospital gardens report positive mood change and reduced stress (Whitehouse et al., 2001). These reports also suggest that gardens and nature in hospitals can heighten patient and family satisfaction with overall quality of care.

***Art in healthcare environments:*** A small number of studies on art in hospitals has yielded findings parallel to those from nature research. Results suggest a consistent pattern wherein the great majority of patients respond positively to representational

nature art, but many react negatively to chaotic abstract art (Ulrich & Gilpin, 2003). For example, Carpman & Grant (1993) studied the preferences of 300 randomly selected inpatients and concluded that the patients consistently preferred nature images but disliked abstract art. Although nature pictures and other emotionally appropriate art elicit positive reactions, there is also evidence that inappropriate art styles or image subject matter can increase stress and worsen other outcomes (Ulrich, 1991). It should not be expected that all art is suitable for high-stress healthcare spaces, as art varies enormously in subject matter and style, and much art is emotionally challenging or provocative.

The pitfalls of displaying emotionally challenging art in healthcare environments are revealed by a study of psychiatric patients (Ulrich, 1991). The unit was extensively furnished with a diverse collection of wall-mounted paintings and prints. Interviews with patients indicated strongly negative reactions to artworks that were ambiguous, surreal, or could be interpreted in multiple ways. The same patients, however, reported having positive feelings and associations with respect to nature paintings and prints.

### ***Provide Social Support***

Many studies of several different categories of patients have indicated that social support reduces stress and improves, for example, recovery outcomes in myocardial infarction patients. Considering the well-established importance of social support, it is unfortunate that there is only a moderate amount of research concerning how hospital design can facilitate or hinder access to social support. Most studies have focused on psychiatric units and nursing homes. There is strong evidence that levels of social interaction can be increased—and presumably beneficial social support as well—by providing lounges, day rooms, and waiting rooms with comfortable movable furniture arranged in small flexible groupings. A few well-designed studies in psychiatric wards and nursing homes have found that appropriate arrangement of movable seating in dining areas enhances social interaction and also improves eating behaviors, such as increasing the amount of food consumed by geriatric patients (Melin & Gotestam, 1981; Peterson, Knapp, Rosen, & al., 1977). Much research on day rooms and waiting areas has shown that the widespread practice of arranging seating side-by-side along room walls inhibits social interaction (Holahan, 1972; Sommer & Ross, 1958). A novel study by Harris (2000) found that family and friends stayed substantially longer during visits to rehabilitation patients when patient rooms were carpeted rather than covered with vinyl flooring.

Much evidence indicates that single rooms are markedly better than multi-bed rooms for supporting or accommodating the presence of family and friends. Some research suggests that open-plan multibed rooms deter family presence and accordingly reduce social support (Sallstrom, Sandman, & Norberg, 1987). Multibed rooms greatly reduce privacy for patient-family interactions compared to single rooms and are much more likely to have restricted visiting hours. A clear advantage of single rooms in fostering social support stems from the fact they provide more space and furniture than double rooms to accommodate family presence (Chaudhury et al., 2003). A survey of staff in four hospitals that each had a mix of single and double rooms found that nurses gave high

ratings to single rooms for accommodating family members but accorded double rooms low scores (Chaudhury et al., 2003). Further, patient-satisfaction data obtained from 2,122,439 patients who received care during 2003 provide overwhelming evidence that patients in single-bed rooms, compared to those with a roommate, are much more satisfied with “accommodations and comfort for family and visitors” (Press Ganey, 2003).

Do patients sharing the same room provide each other with stress-reducing social support? While some patients find roommates provide comforting social support, findings from several studies indicate that the presence of a roommate usually is a source of stress rather than social support. In most cases, roommates are linked to stressors, for example, loss of privacy or having a roommate who is unfriendly, has too many visitors, or is seriously ill (Van der Ploeg, 1988; Volicer, Isenberg, & Burns, 1977). An earlier section emphasized that noise is a much greater problem in double rooms than singles, and that noises stemming from the presence of other patients are a major cause of sleep loss.

### ***Improve Communication to Patients***

Good staff communication helps reduce patient and family anxiety, promotes better care at home after discharge, and in other ways can improve outcomes. Good communication also tends to be the single most important factor affecting overall satisfaction with care across different patient categories (Press Ganey, 2003). Data obtained from 2,122,439 patients nationally in 2003 show that patients consistently report significantly higher satisfaction with communication from nurses and physicians when they are in single rooms compared to when they have one or more roommates (Press Ganey, 2003). To explain this clear and important advantage of single rooms, Kaldenburg (1999) has proposed that staff in multibed rooms are reluctant to discuss patient issues or give information within hearing of a roommate, out of respect for privacy. Growing concern for patient confidentiality and HIPAA are certain to increase the already major advantages of single rooms with respect to communication.

## **IV. Improve Overall Healthcare Quality**

### ***Provide Single-Bed Patient Rooms***

Based on an extremely large and varied body of research reviewed in earlier sections, there can be no question that single-bed rooms have several major advantages over double rooms and open bays. To summarize briefly, these advantages include: lower nosocomial infection rates, fewer patient transfers and associated medical errors, far less noise, much better patient privacy and confidentiality, better communication from staff to patients and from patients to staff, superior accommodation of family and consistently higher satisfaction with overall quality of care.



### ***Reduce Length of Stay***

Climate and sunlight influences length of hospital stay as well as sleep-wake patterns among hospitalized patients (Beauchemin & Hays, 1996; Benedetti, Francesco, Colombo, Cristina, Barbini, Barbara, Campori, Euridice, & Smeraldi, Enrico, 2001; Federman, Drebing, Boisvert, Penk, 2000; Hebert, Dumont, & Paquet, 1998; Kecskes et al., 2003; Kinnunen, Saynajakangas, Tuuponen, & Keistinen, 2002). One research group studied the impact of the amount of natural light on the length of hospitalization of patients with unipolar and bipolar disorder. The researchers found that bipolar patients randomly assigned to the brighter, eastern rooms (exposed to direct sunlight in the morning) had a mean 3.67-day shorter hospital stay than patients in west-facing rooms (Benedetti, Francesco et al., 2001). Patients recovering from abdominal surgery had shorter stays if they had a bedside window view of nature rather than if their windows looked out onto a brick wall (Ulrich, 1984).

The large research literature on infection reviewed in an earlier section indicated that the design of the physical environment strongly impacts hospital-acquired infection rates by affecting both airborne and contact transmission routes. Evidence-based design measures, by reducing nosocomial infection rates, play a key role in shortening hospital stays.

### ***Increase Patient Satisfaction with Quality of Care***

There is strong evidence that design changes that make the environment more comfortable, aesthetically pleasing, and informative relieve stress among patients and increases satisfaction with the quality of care provided. Renovating a traditional waiting area in a neurology clinic by making small changes to the general layout, color scheme, furniture, floor covering, curtains, and providing informational material and information displays resulted in more positive environmental appraisals, improved mood, altered physiological state, and greater reported satisfaction among waiting patients (Leather, Beale, Santos, Watts, & Lee, 2003).

Patients in well-decorated and well-appointed hotel-like rooms rated their attending physicians, housekeeping, and food-service staff, the food, and the hospital better than patients in standard rooms (typical hospital beds, inexpensive family sitting chairs, and no artwork) in the same hospital. Also, they had stronger intentions to use the hospital again and would recommend the hospital to others (Swan, Richardson, & Hutton, 2003). In another study, it was found that environmental satisfaction was a significant predictor of overall satisfaction, ranking only below perceived quality of nursing and clinical care (Harris, P. B., McBride, Ross, & Curtis, 2002). Several post-occupancy evaluation studies have examined patient and staff satisfaction with the different elements of the healthcare environment such as gardens, individual wards, and patient rooms (Brown, Wright, & Brown, 1997; Heath & Gifford, 2001; Shepley, 1995, 2002; Shepley & Davies, 2003; Shepley & Wilson, 1999; Whitehouse et al., 2001). These studies yield rich context-specific data that describe which aspects of the environment were effective. Some studies show quite clearly that tacit, as well implicit, staff and organizational practices and policies influence how an environment actually functions and is perceived

by patients and staff members. The qualitative and quantitative data provided by such studies are very important in understanding the nature of the problem (e.g. why are the gardens not being used as intended?) as well in developing solutions to tackle the problem. Future research that looks at satisfaction among hospital patients should consider using these multi-method post-occupancy evaluations that use different methods to obtain objective and subjective evaluations of use and satisfaction.

## CONCLUSIONS

The research team found more than 600 rigorous studies linking a range of aspects of the built environment of hospitals to staff stress and effectiveness, patient safety, patient and family stress and healing, and improved overall healthcare quality and cost. This deep and wide base of evidence suggests that, parallel to evidence-based medicine, we can move to evidence-based design (EBD). EBD refers to a process for creating healthcare buildings, informed by the best available evidence, with the goal of improving outcomes and of continuing to monitor the success of designs for subsequent decision-making. While it is difficult to conduct rigorous research on the impacts of the healthcare environment—hospitals are complex systems where it is difficult to isolate the impact of single factors and the building industry conducts little impact-based research—The Center for Health Design and other groups have made considerable progress in developing a knowledge base of evidence.

EBD is not about hospitals that are simply nicer or fancier than traditional hospitals. Rather, the focus of evidence-based design is to create hospitals that actually help patients recover and be safer, and help staff do their jobs better. EBD is a process for creating health care buildings informed by the best available evidence concerning how the physical environment can interfere with or support activities by patients, families, and staff, and how the setting provides experiences that provide a caring, effective, safe, patient-centered environment. Many of the improvements suggested by EBD are only slightly more expensive than traditional solutions, if they are more expensive at all.

The large research literature surveyed in this report point to several actions we can take immediately:

- Provide single-bed rooms in almost all situations. Adaptable-acuity single-bed rooms should be widely adopted. Single rooms have been shown to lower hospital-induced nosocomial infections, reduce room transfers and associated medical errors, greatly lessen noise, improve patient confidentiality and privacy, facilitate social support by families, improve staff communication to patients, and increase patients' overall satisfaction with health care.
- New hospitals should be much quieter to reduce stress and improve sleep and other outcomes. Noise levels will be substantially lowered by the following combination of environmental interventions: providing single-bed rooms, installing high-performance sound-absorbing ceilings, and eliminating noise sources (for example, using noiseless paging).

- Provide patients stress reducing views of nature and other positive distractions.
- Develop wayfinding systems that allow users, and particularly outpatients and visitors, to find their way efficiently and with little stress.
- Improve ventilation through the use of improved filters, attention to appropriate pressurization, and special vigilance during construction.
- Improve lighting, especially access to natural lighting and full-spectrum lighting.
- Design ward layouts and nurses stations to reduce staff walking and fatigue, increase patient care time, and support staff activities such as medication supply, communication, charting, and respite from stress.

## REFERENCES CITED

- Aaron, J. N., Carlisle, C. C., Carskadon, M. A., Meyer, T. J., Hill, N. S., & Millman, R. P. (1996). Environmental noise as a cause of sleep disruption in an intermediate respiratory care unit. *Sleep*, *19*(9), 707-710.
- Albert, R. K., & Condie, F. (1981). Hand-washing patterns in medical intensive-care units. *New England Journal of Medicine*, *304*(24), 1465-1466.
- Alberti, C., Bouakline, A., Ribaud, P., Lacroix, C., Rousselot, P., Leblanc, T., et al. (2001). Relationship between environmental fungal contamination and the incidence of invasive aspergillosis in haematology patients. *The Journal of Hospital Infection*, *48*(3), 198-206.
- Allaouchiche, B., Dufflo, F., Debon, R., Bergeret, A., & Chassard, D. (2002). Noise in the postanaesthesia care unit. *British Journal of Anaesthesia*, *88*(3), 369-373.
- Archibald, L. K., Manning, M. L., Bell, L. M., Banerjee, S., & Jarvis, W. R. (1997). Patient density, nurse-to-patient ratio and nosocomial infection risk in a pediatric cardiac intensive care unit. *The Pediatric Infectious Disease Journal*, *16*(11), 1045-1048.
- Arlet, G., Gluckman, E., Gerber, F., Perol, Y., & Hirsch, A. (1989). Measurement of bacterial and fungal air counts in two bone marrow transplant units. *The Journal of Hospital Infection*, *13*(1), 63-69.
- Aygun, G., Demirkiran, O., Utku, T., Mete, B., Urkmez, S., Yilmaz, M., et al. (2002). Environmental contamination during a carbapenem-resistant *Acinetobacter baumannii* outbreak in an intensive care unit. *The Journal of Hospital Infection*, *52*(4), 259-262.
- Babwin, D. 2002. Building Boom. *Hospitals & Health Networks*. *76*(3): 48-54
- Baker, C. F. (1984). Sensory overload and noise in the ICU: Sources of environmental stress. *Critical Care Quarterly*, *6*(4), 66-80.
- Baker, C. F. (1992). Discomfort to environmental noise: Heart rate responses of SICU patients. *Critical Care Nursing Quarterly*, *15*(2), 75-90.
- Barlas, D., Sama, A. E., Ward, M. F., & Lesser, M. L. (2001). Comparison of the auditory and visual privacy of emergency department treatment areas with curtains versus those with solid walls. *Annals of Emergency Medicine*, *38*(2), 135-139.
- Bauer, T. M., Ofner, E., Just, H. M., Just, H., & Daschner, F. D. (1990). An epidemiological study assessing the relative importance of airborne and direct contact transmission of microorganisms in a medical intensive care unit. *The Journal of Hospital Infection*, *15*(4), 301-309.
- Bayo, M. V., Garcia, A. M., & Garcia, A. (1995). Noise levels in an urban hospital and workers' subjective responses. *Archives of Environmental Health*, *50*(3), 247-251.
- Beauchemin, K. M., & Hays, P. (1996). Sunny hospital rooms expedite recovery from severe and refractory depressions. *Journal of Affective Disorders*, *40*(1-2), 49-51.
- Ben-Abraham, R., Keller, N., Szold, O., Vardi, A., Weinberg, M., Barzilay, Z., et al. (2002). Do isolation rooms reduce the rate of nosocomial infections in the pediatric intensive care unit? *Journal of Critical Care*, *17*(3), 176-180.
- Benedetti, F., Colombo, C., Barbini, B., Campori, E., & Smeraldi, E. (2001). Morning sunlight reduces length of hospitalization in bipolar depression. *Journal of Affective Disorders*, *62*(3), 221-223.
- Berg, S. (2001). Impact of reduced reverberation time on sound-induced arousals during sleep. *Sleep*, *24*(3), 289-292.
- Berglund, B., Lindvall, T., & Schwela, D. H. (1999). *Guidelines for community noise*. World Health Organization: Protection of the Human Environment.
- Blomkvist, V., Eriksen, C. A., Theorell, T., Ulrich, R. S., & Rasmanis, G. (in press, 2004). Acoustics and psychosocial environment in coronary intensive care. *Occupational and Environmental Medicine*.
- Booker, J. M., & Roseman, C. (1995). A seasonal pattern of hospital medication errors in Alaska. *Psychiatry Research*, *57*(3), 251-257.
- Boyce, J. M., Potter-Bynoe, G., Chenevert, C., & King, T. (1997). Environmental contamination due to methicillin-resistant *Staphylococcus aureus*: Possible infection control implications. *Infection Control and Hospital Epidemiology*, *18*(9), 622-627.
- Brandis, S. (1999). A collaborative occupational therapy and nursing approach to falls prevention in hospital inpatients. *Journal of Quality in Clinical Practice*, *19*(4), 215-221.

- Brown, B., Wright, H., & Brown, C. (1997). A post-occupancy evaluation of wayfinding in a pediatric hospital: Research findings and implications for instruction. *Journal of Architectural & Planning Research, 14*(1), 35-51.
- Buchanan, T. L., Barker, K. N., Gibson, J. T., Jiang, B. C., & Pearson, R. E. (1991). Illumination and errors in dispensing. *American Journal of Hospital Pharmacy, 48*(10), 2137-2145.
- Bures, S., Fishbain, J. T., Uyehara, C. F., Parker, J. M., & Berg, B. W. (2000). Computer keyboards and faucet handles as reservoirs of nosocomial pathogens in the intensive care unit. *American Journal of Infection Control, 28*(6), 465-471.
- Burgio, L., Engel, B., Hawkins, A., McCorick, K., & Scheve, A. (1990). A descriptive analysis of nursing staff behaviors in a teaching nursing home: Differences among NAs, LPNs and RNs. *The Gerontologist, 30*, 107-112.
- Butler, D., Acquino, A. L., Hissong, A. A., & Scott, P. A. (1993). Wayfinding by newcomers in a complex building. *Human Factors, 25*(1), 159-173.
- Capezuti, E., Maislin, G., Strumpf, N., & Evans, L. K. (2002). Side rail use and bed-related fall outcomes among nursing home residents. *Journal of the American Geriatrics Society, 50*(1), 90-96.
- Carpman, J., & Grant, M. (1993). *Design that cares: Planning health facilities for patients and visitors* (2nd ed.). Chicago: American Hospital Publishing.
- Carpman, J., Grant, M., & Simmons, D. (1983-84). Wayfinding in the hospital environment: The impact of various floor numbering alternatives. *Journal of Environmental Systems, 13*(4), 353-364.
- Carpman, J., Grant, M., & Simmons, D. (1984). *No more mazes: Research about design for wayfinding in hospitals*. Ann Arbor, Michigan: The University of Michigan Hospitals.
- Carpman, J., Grant, M. A., & Simmons, D. A. (1985). Hospital design and wayfinding: A video simulation study. *Environment & Behavior, 17*(3), 296-314.
- Chang, J. T., Morton, S. C., Rubenstein, L. Z., Mojica, W. A., Maglione, M., Suttorp, M. J., et al. (2004). Interventions for the prevention of falls in older adults: Systematic review and meta-analysis of randomised clinical trials. *British Medical Journal, 328*(7441), 680-680.
- Chang, V. T., & Nelson, K. (2000). The role of physical proximity in nosocomial diarrhea. *Clinical Infectious Diseases, 31*(3), 717-722.
- Chaudhury, H., Mahmood, A., & Valente, M. (2003). *Pilot study on comparative assessment of patient care issues in single and multiple occupancy rooms* (Unpublished report): The Coalition for Health Environments Research.
- Christensen, K. E. (1979). *An impact analysis framework for calculating the costs of staff disorientation in hospitals*. Los Angeles: School of Architecture and Urban Planning, University of California.
- Cmiel, C. A., Karr, D. M., Gasser, D. M., Oliphant, L. M., & Neveau, A. J. (2004). Noise control: A nursing team's approach to sleep promotion. *American Journal of Nursing, 104*(2), 40-48.
- Cohen, B., Saiman, L., Cimiotti, J., & Larson, E. (2003). Factors associated with hand hygiene practices in two neonatal intensive care units. *The Pediatric Infectious Diseases Journal, 22*(6), 494-499.
- Conly, J. M., Hill, S., Ross, J., Lertzman, J., & Louie, T. J. (1989). Handwashing practices in an intensive care unit: The effects of an educational program and its relationship to infection rates. *American Journal of Infection Control, 17*(6), 330-339.
- Cook, R. I., Render, M., & Woods, D. D. (2000). Gaps in the continuity of care and progress on patient safety. *British Medical Journal, 320*(7), 791-794.
- Cooper Marcus, C., & Barnes, M. (1995). *Gardens in healthcare facilities: Uses, therapeutic benefits, and design recommendations*. Martinez, CA: Center for Health Design.
- Cornet, M., Levy, V., Fleury, L., Lortholary, J., Barquins, S., Coureul, M. H., et al. (1999). Efficacy of prevention by high-efficiency particulate air filtration or laminar airflow against *Aspergillus* airborne contamination during hospital renovation. *Infection Control and Hospital Epidemiology, 20*(7), 508-513.
- Couper, R. T., Hendy, K., Lloyd, N., Gray, N., Williams, S., & Bates, D. J. (1994). Traffic and noise in children's wards. *Medical Journal of Australia, 160*(6), 338-341.
- Devine, J., Cooke, R. P., & Wright, E. P. (2001). Is methicillin-resistant *Staphylococcus aureus* (MRSA) contamination of ward-based computer terminals a surrogate marker for nosocomial MRSA transmission and handwashing compliance? *The Journal of Hospital Infection, 48*(1), 72-75.
- Dharan, S., & Pittet, D. (2002). Environmental controls in operating theatres. *The Journal of Hospital Infection, 51*(2), 79-84.

- Diette, G. B., Lechtzin, N., Haponik, E., Devrotes, A., & Rubin, H. R. (2003). Distraction therapy with nature sights and sounds reduces pain during flexible bronchoscopy: A complementary approach to routine analgesia. *Chest*, *123*(3), 941-948.
- Dorsey, S. T., Cydulka, R. K., & Emerman, C. L. (1996). Is handwashing teachable? Failure to improve handwashing behavior in an urban emergency department. *Academy of Emergency Medicine*, *3*(4), 360-365.
- Dubbert, P. M., Dolce, J., Richter, W., Miller, M., & Chapman, S. W. (1990). Increasing ICU staff handwashing: Effects of education and group feedback. *Infection Control and Hospital Epidemiology*, *11*(4), 191-193.
- Falk, S. A., & Woods, N. F. (1973). Hospital noise: Levels and potential health hazards. *New England Journal of Medicine*, *289*(15), 774-781.
- Farquharson, C., & Baguley, K. (2003). Responding to the severe acute respiratory syndrome (SARS) outbreak: Lessons learned in a Toronto emergency department. *Journal of Emergency Nursing*, *29*(3), 222-228.
- Federman, E. J., Drebing, C. E., Boisvert, C., Penk, W., Binus, G., & Rosenheck, R. (2000). Relationship between climate and psychiatric inpatient length of stay in Veterans Health Administration hospitals. *American Journal of Psychiatry*, *157*(10), 1669.
- Flynn, E. A., Barker, K. N., Gibson, J. T., Pearson, R. E., Berger, B. A., & Smith, L. A. (1999). Impact of interruptions and distractions on dispensing errors in an ambulatory care pharmacy. *American Journal of Health Systems Pharmacy*, *56*(13), 1319-1325.
- Fox, R. A., & Henson, P. W. (1996). Potential ocular hazard from a surgical light source. *Australasian Physical and Engineering Sciences in Medicine*, *19*(1), 12-16.
- Friberg, S., Ardnor, B., & Lundholm, R. (2003). The addition of a mobile ultra-clean exponential laminar airflow screen to conventional operating room ventilation reduces bacterial contamination to operating box levels. *The Journal of Hospital Infection*, *55*(2), 92-97.
- Gabor, J. Y., Cooper, A. B., Crombach, S. A., Lee, B., Kadikar, N., Bettger, H. E., et al. (2003). Contribution of the intensive care unit environment to sleep disruption in mechanically ventilated patients and healthy subjects. *American Journal of Respiratory and Critical Care Medicine*, *167*(5), 708-715.
- Gardner, P. S., Court, S. D., Brocklebank, J. T., Downham, M. A., & Weightman, D. (1973). Virus cross-infection in paediatric wards. *British Medical Journal*, *2*(5866), 571-575.
- Garg, A., & Owen, B. (1992). Reducing back stress to nursing personnel: An ergonomic intervention in a nursing home. *Ergonomics*, *35*(11), 1353-1375.
- Gast, P. L., & Baker, C. F. (1989). The CCU patient: Anxiety and annoyance to noise. *Critical Care Nursing Quarterly*, *12*(3), 39-54.
- Goldmann, D. A., Durbin, W. A., Jr., & Freeman, J. (1981). Nosocomial infections in a neonatal intensive care unit. *Journal of Infectious Diseases*, *144*(5), 449-459.
- Gotlieb, J. B. (2000). Understanding the effects of nurses, patients' hospital rooms, and patients' perception of control in the perceived quality of a hospital. *Health Marketing Quarterly*, *18*(1/2), 1.
- Graham, M. (1990). Frequency and duration of handwashing in an intensive care unit. *American Journal of Infection Control*, *18*(2), 77-81.
- Grover, P. (1971). *Wayfinding in hospital environments: UCLA hospital disorientation pilot case study*. Los Angeles, Calif.: Graduate School of Architecture and Urban Planning, University of California, Los Angeles.
- Hagerman, I., Theorell, T., Ulrich, R. S., Blomkvist, V., Eriksen, C. A., & Rasmanis, G. (in press, 2004). Influence of coronary intensive care acoustics on the physiological states and quality of care of patients. *International Journal of Cardiology*.
- Hahn, T., Cummings, K. M., Michalek, A. M., Lipman, B. J., Segal, B. H., & McCarthy, P. L., Jr. (2002). Efficacy of high-efficiency particulate air filtration in preventing aspergillosis in immunocompromised patients with hematologic malignancies. *Infection Control and Hospital Epidemiology*, *23*(9), 525-531.
- Hamilton, K. (2003). The four levels of evidence based practice. *Healthcare Design*, *3*, 18-26.
- Hanger, H. C., Ball, M. C., & Wood, L. A. (1999). An analysis of falls in the hospital: Can we do without bedrails? *Journal of the American Geriatrics Society*, *47*(5), 529-531.

- Haq, S., & Zimring, C. (2003). Just down the road a piece: The development of topological knowledge of building layouts. *Environment & Behavior*, 35(1), 132-160.
- Harris, D. (2000). *Environmental quality and healing environments: A study of flooring materials in a healthcare telemetry unit*. Doctoral dissertation, Texas A&M University, College Station.
- Harris, P. B., McBride, G., Ross, C., & Curtis, L. (2002). A place to heal: Environmental sources of satisfaction among hospital patients. *Journal of Applied Social Psychology*, 32(6), 1276-1299.
- Heath, Y., & Gifford, R. (2001). Post-occupancy evaluation of therapeutic gardens in a multi-level care facility for the aged. *Activities, Adaptation & Aging*, 25(2), 21-43.
- Hebert, M., Dumont, M., & Paquet, J. (1998). Seasonal and diurnal patterns of human illumination under natural conditions. *Chronobiology International*, 15(1), 59-70.
- Hendrich, A. (2003). *Optimizing physical space for improved outcomes: Satisfaction and the bottom line*. Paper presented at "Optimizing the Physical Space for Improved Outcomes, Satisfaction, and the Bottom Line," minicourse sponsored by the Institute for Healthcare Improvement and the Center for Health Design, Atlanta, GA.
- Hendrich, A., Fay, J., & Sorrells, A. (2002, September). Courage to heal: Comprehensive cardiac critical care. *Healthcare Design*, 11-13
- Hendrich, A., Fay, J., and Sorrells, A. (2004). Effects of acuity-adaptable rooms on flow of patients and delivery of care. *American Journal of Critical Care*, 13 (1).
- Hibbard, F. J., Bosso, J. A., Sward, L. W., & Baum, S. (1981). Delivery time in a decentralized pharmacy system without satellites. *American Journal of Hospital Pharmacy*, 38(5), 690-692.
- Hilton, B. A. (1985). Noise in acute patient care areas. *Research in Nursing & Health*, 8(3), 283-291.
- Hodge, B., & Thompson, J. F. (1990). Noise pollution in the operating theatre. *Lancet*, 335(8694), 891-894.
- Holahan, C.J. (1972). Seating patterns and patient behavior in an experimental dayroom. *Journal of Abnormal Psychology*, 80(2), 115-124.
- Humphreys, H., Johnson, E. M., Warnock, D. W., Willatts, S. M., Winter, R. J., & Speller, D. C. (1991). An outbreak of aspergillosis in a general ITU. *The Journal of Hospital Infection*, 18(3), 167-177.
- Institute of Medicine (IOM). (2004). *Keeping patients safe: Transforming the work environment of nurses*. Washington, DC: National Academy Press.
- Institute of Medicine (IOM) (2001). *Crossing the Quality Chasm: A new health system for the 21st century*. Washington, DC: National Academy Press.
- Iwen, P. C., Davis, J. C., Reed, E. C., Winfield, B. A., & Hinrichs, S. H. (1994). Airborne fungal spore monitoring in a protective environment during hospital construction, and correlation with an outbreak of invasive aspergillosis. *Infection Control and Hospital Epidemiology*, 15(5), 303-306.
- Joint Commission on Accreditation of Healthcare Organizations (2002). *Health care at the crossroad: Strategies for addressing the evolving nursing crisis*. Oakbrook Terrace, IL: Joint Commission on Accreditation of Healthcare Organizations.
- Jernigan, J. A., Titus, M. G., Groschel, D. H., Getchell-White, S., & Farr, B. M. (1996). Effectiveness of contact isolation during a hospital outbreak of methicillin-resistant *Staphylococcus aureus*. *American Journal of Epidemiology*, 143(5), 496-504.
- Jiang, S., Huang, L., Chen, X., Wang, J., Wu, W., Yin, S., et al. (2003). Ventilation of wards and nosocomial outbreak of severe acute respiratory syndrome among healthcare workers. *Chinese Medical Journal*, 116(9), 1293-1297.
- Johnson, A. N. (2001). Neonatal response to control of noise inside the incubator. *Pediatric Nursing*, 27(6), 600-605.
- Kaldenburg, D. O. (1999). The influence of having a roommate on patient satisfaction. *Satisfaction Monitor*, January-February (www.pressganey.org).
- Kaplan, L. M., & McGuckin, M. (1986). Increasing handwashing compliance with more accessible sinks. *Infection Control*, 7(8), 408-410.
- Kecskes, I., Rihmer, Z., Kiss, K., Vargha, A., Szili, I., & Rihmer, A. (2003). Possible effect of gender and season on the length of hospitalisation in unipolar major depressives. *Journal of Affective Disorders*, 73(3), 279-282.
- Kinnunen, T., Saynajakangas, O., Tuuponen, T., & Keistinen, T. (2002). Regional and seasonal variation in the length of hospital stay for chronic obstructive pulmonary disease in Finland. *International Journal of Circumpolar Health*, 61(2), 131-135.

- Kistner, U. A., Keith, M. R., Sergeant, K. A., & Hokanson, J. A. (1994). Accuracy of dispensing in a high-volume, hospital-based outpatient pharmacy. *American Journal of Hospital Pharmacy*, 51(22), 2793-2797.
- Kromhout, H., Hoek, F., Uitterhoeve, R., Huijbers, R., Overmars, R. F., Anzion, R., et al. (2000). Postulating a dermal pathway for exposure to anti-neoplastic drugs among hospital workers: Applying a conceptual model to the results of three workplace surveys. *The Annals of Occupational Hygiene*, 44(7), 551-560.
- Kumari, D. N., Haji, T. C., Keer, V., Hawkey, P. M., Duncanson, V., & Flower, E. (1998). Ventilation grilles as a potential source of methicillin-resistant *Staphylococcus aureus* causing an outbreak in an orthopaedic ward at a district general hospital. *The Journal of Hospital Infection*, 39(2), 127-133.
- Larson, E. (1988). A causal link between handwashing and risk of infection? Examination of the evidence. *Infection Control*, 9(1), 28-36.
- Larson, E., McGeer, A., Quraishi, Z. A., Krenzschek, D., Parsons, B. J., Holdford, J., et al. (1991). Effect of an automated sink on handwashing practices and attitudes in high-risk units. *Infection Control and Hospital Epidemiology*, 12(7), 422-428.
- Larson, E. L., Bryan, J. L., Adler, L. M., & Blane, C. (1997). A multifaceted approach to changing handwashing behavior. *American Journal of Infection Control*, 25(1), 3-10.
- Leather, P., Beale, D., Santos, A., Watts, J., & Lee, L. (2003). Outcomes of environmental appraisal of different hospital waiting areas. *Environment and Behavior*, 35(6), 842-869.
- Levine, M., Marchon, I., & Hanley, G. (1984). The placement and misplacement of you-are-here maps. *Environment & Behavior*, 16(2), 139-157.
- Lewy, A. J., Bauer, V. K., Cutler, N. L., Sack, R. L., Ahmed, S., Thomas, K. H., et al. (1998). Morning vs evening light treatment of patients with winter depression. *Archives of General Psychiatry*, 55(10), 890-896.
- Loo, V. G., Bertrand, C., Dixon, C., Vitye, D., DeSalis, B., McLean, A. P., et al. (1996). Control of construction-associated nosocomial aspergillosis in an antiquated hematology unit. *Infection Control and Hospital Epidemiology*, 17(6), 360-364.
- Love, H. (2003). Noise exposure in the orthopaedic operating theatre: A significant health hazard. *ANZ Journal of Surgery*, 73(10), 836-838.
- Lovell, B. B., Ancoli-Israel, S., & Gevirtz, R. (1995). Effect of bright light treatment on agitated behavior in institutionalized elderly subjects. *Psychiatry Research*, 57(1), 7-12.
- Lutz, B. D. J., Jiankang Rinaldi, Michael G Wickes, Brian L Huycke, Mark M. (2003). Outbreak of invasive Aspergillus infection in surgical patients, associated with a contaminated air-handling system. *Clinical Infectious Diseases*, 37(6), 786-793.
- Mahieu, L. M., De Dooy, J. J., Van Laer, F. A., Jansens, H., & Ieven, M. M. (2000). A prospective study on factors influencing aspergillus spore load in the air during renovation works in a neonatal intensive care unit. *The Journal of Hospital Infection*, 45(3), 191-197.
- McDonald, L. C., Walker, M., Carson, L., Arduino, M., Aguero, S. M., Gomez, P., et al. (1998). Outbreak of *Acinetobacter* spp. bloodstream infections in a nursery associated with contaminated aerosols and air conditioners. *The Pediatric Infectious Disease Journal*, 17(8), 716-722.
- McKendrick, G. D., & Emond, R. T. (1976). Investigation of cross-infection in isolation wards of different design. *Journal of Hygiene (Lond)*, 76(1), 23-31.
- McLaughlin, A., McLaughlin, B., Elliott, J., & Campalani, G. (1996). Noise levels in a cardiac surgical intensive care unit: A preliminary study conducted in secret. *Intensive Critical Care Nursing*, 12(4), 226-230.
- McManus, A. T., Mason, A. D., Jr., McManus, W. F., & Pruitt, B. A., Jr. (1994). A decade of reduced gram-negative infections and mortality associated with improved isolation of burned patients. *Archives of Surgery*, 129(12), 1306-1309.
- McManus, A. T., Mason, A. D. Jr., McManus, W. F., & Pruitt, B. A. Jr. (1992). Control of pseudomonas aeruginosa infections in burned patients. *Surgical Research Communications*, 12, 61-67.
- McManus, A. T., McManus, W. F., Mason, A. D., Jr., Aitchison, A. R., & Pruitt, B. A., Jr. (1985). Microbial colonization in a new intensive care burn unit. A prospective cohort study. *Archives of Surgery*, 120(2), 217-223.



- Melin, L., & Gotestam, K. G. (1981). The effects of rearranging ward routines on communication and eating behaviors of psychogeriatric patients. *Journal of Applied Behavior Analysis, 14*(1), 47-51.
- Menzies, D., Fanning, A., Yuan, L., & FitzGerald, J. M. (2000). Hospital ventilation and risk for tuberculosis infection in Canadian health care workers. *Annals of Internal Medicine, 133*(10), 779-789.
- Meyer, T. J., Eveloff, S. E., Bauer, M. S., Schwartz, W. A., Hill, N. S., & Millman, R. P. (1994). Adverse environmental conditions in the respiratory and medical ICU settings. *Chest, 105*(4), 1211-1216.
- Miller, A. C., Hickman, L. C., & Lemasters, G. K. (1992). A distraction technique for control of burn pain. *Journal of Burn Care and Rehabilitation, 13*(5), 576-580.
- Mlinek, E. J., & Pierce, J. (1997). Confidentiality and privacy breaches in a university hospital emergency department. *Academy of Emergency Medicine, 4*(12), 1142-1146.
- Moeser, S. D. (1988). Cognitive mapping in a complex building. *Environment & Behavior, 20*(1), 21-49.
- Moore, M. M. N., Stanton P. Nguyen, Diem Robinson, Samuel P. Ryals, Brenda Imbrie, John Z. Spotnitz, William. (1998). Interventions to reduce decibel levels on patient care units, *American Surgeon, 64*(9), 894.
- Morrison, W. E., Haas, E. C., Shaffner, D. H., Garrett, E. S., & Fackler, J. C. (2003). Noise, stress, and annoyance in a pediatric intensive care unit. *Critical Care Medicine, 31*(1), 113-119.
- Mulin, B., Rouget, C., Clement, C., Bailly, P., Julliot, M. C., Viel, J. F., et al. (1997). Association of private isolation rooms with ventilator-associated *Acinetobacter baumannii* pneumonia in a surgical intensive-care unit. *Infection Control and Hospital Epidemiology, 18*(7), 499-503.
- Muto, C. A., Siström, M. G., & Farr, B. M. (2000). Hand hygiene rates unaffected by installation of dispensers of a rapidly acting hand antiseptic. *American Journal of Infection Control, 28*(3), 273-276.
- Neely, A. N., & Maley, M. P. (2001). Dealing with contaminated computer keyboards and microbial survival. *American Journal of Infection Control, 29*(2), 131-132.
- Nelson-Shulman, Y. (1983-84). Information and environmental stress: Report of a hospital intervention. *Journal of Environmental Systems, 13*(4), 303-316.
- Norbeck, J. S. (1985). Perceived job stress, job satisfaction, and psychological symptoms in critical care nursing. *Research in Nursing & Health, 8*(3), 253-259.
- Noskin, G. A., Bednarz, P., Suriano, T., Reiner, S., & Peterson, L. (2000). Persistent contamination of fabric-covered furniture by Vancomycin-resistant Enterococci: Implication for upholstery selection in hospitals. *American Journal of Infection Control, 28*(4), 311-313.
- Nott, M. R., & West, P. D. (2003). Orthopaedic theatre noise: A potential hazard to patients. *Anaesthesia, 58*(8), 784-787.
- Novaes, M. A., Aronovich, A., Ferraz, M. B., & Knobel, E. (1997). Stressors in ICU: patients' evaluation. *Intensive Care Medicine, 23*(12), 1282-1285.
- Opal, S. M., Asp, A. A., Cannady, P. B., Jr., Morse, P. L., Burton, L. J., & Hammer, P. G. (1986). Efficacy of infection control measures during a nosocomial outbreak of disseminated aspergillosis associated with hospital construction. *Journal of Infectious Diseases, 153*(3), 634-637.
- Oren, I., Haddad, N., Finkelstein, R., & Rowe, J. M. (2001). Invasive pulmonary aspergillosis in neutropenic patients during hospital construction: Before and after chemoprophylaxis and institution of HEPA filters. *American Journal of Hematology, 66*(4), 257-262.
- Ortega-Andeane, P., & Urbina-Soria, J. (1988). *A case study of wayfinding and security in a Mexico City hospital*. Paper presented at the EDRA: Environmental Design Research Association, US, 19:231-236.
- Palmer, R. (1999). Bacterial contamination of curtains in clinical areas. *Nursing Standard, 14*(2), 33-35.
- Parsons, R., & Hartig, T. (2000). Environmental psychophysiology. In J. T. Cacioppo & L. G. Tassinary (Eds.), *Handbook of psychophysiology (2nd ed.)* (pp. 815-846). New York: Cambridge University Press.
- Parthasarathy, S., & Tobin, M. J. (2004). Sleep in the intensive care unit. *Intensive Care Medicine, 30*(2), 197-206.
- Passini, R., Rainville, C., Marchand, N., & Joannette, Y. (1995). Wayfinding in dementia of the Alzheimer type: Planning abilities. *Journal of Clinical and Experimental Neuropsychology, 17*(6), 820-832.

- Passweg, J. R., Rowlings, P. A., Atkinson, K. A., Barrett, A. J., Gale, R. P., Gratwohl, A., et al. (1998). Influence of protective isolation on outcome of allogeneic bone marrow transplantation for leukemia. *Bone Marrow Transplant*, *21*(12), 1231-1238.
- Pegues, D. A., & Woernle, C. H. (1993). An outbreak of acute nonbacterial gastroenteritis in a nursing home. *Infection Control and Hospital Epidemiology*, *14*(2), 87-94.
- Peponis, J., Zimring, C., & Choi, Y. K. (1990). Finding the building in wayfinding. *Environment & Behavior*, *22*(5), 555-590.
- Peterson, R., Knapp, T., Rosen, J., & Pither, B.F. (1977). The effects of furniture arrangement on the behavior of geriatric patients. *Behavior Therapy*, *8*, 464-467.
- Pierce, R. A., 2nd, Rogers, E. M., Sharp, M. H., & Musulin, M. (1990). Outpatient pharmacy redesign to improve work flow, waiting time, and patient satisfaction. *American Journal of Hospital Pharmacy*, *47*(2), 351-356.
- Pittet, D., Hugonnet, S., Harbarth, S., Mourouga, P., Sauvan, V., Touveneau, S., et al. (2000). Effectiveness of a hospital-wide programme to improve compliance with hand hygiene. *Lancet*, *356*(9238), 1307-1312.
- Press Ganey, Inc. (2003). National patient satisfaction data for 2003. (Provided by Press Ganey, Inc. at the request of the authors for this research review.)
- Preston, G. A., Larson, E. L., & Stamm, W. E. (1981). The effect of private isolation rooms on patient care practices, Colonization and infection in an intensive care unit. *The American Journal of Medicine*, *70*(3), 641-645.
- Reynolds, D. M., Johnson, M. H., & Longe, R. L. (1978). Medication delivery time requirements in centralized and decentralized unit dose drug distribution systems. *American Journal of Hospital Pharmacy*, *35*(8), 941-943.
- Roberts, S. A., Findlay, R., & Lang, S. D. (2001). Investigation of an outbreak of multi-drug resistant *Acinetobacter baumannii* in an intensive care burns unit. *The Journal of Hospital Infection*, *48*(3), 228-232.
- Robertson, A., Cooper-Peel, C., & Vos, P. (1998). Peak noise distribution in the neonatal intensive care nursery. *Journal of Perinatology*, *18*(5), 361-364.
- Rountree, P. M., Beard, M. A., Loewenthal, J., May, J., & Renwick, S. B. (1967). Staphylococcal sepsis in a new surgical ward. *British Medical Journal*, *1*(533), 132-137.
- Rubin, H., Owens, A. J., & Golden, G. (1998). *Status report (1998): An investigation to determine whether the built environment affects patients' medical outcomes*. Martinez, CA: Center for Health Design.
- Sallstrom, C., Sandman, P. O., & Norberg, A. (1987). Relatives' experience of the terminal care of long-term geriatric patients in open-plan rooms. *Scandinavian Journal of Caring Science*, *1*(4), 133-140.
- Sanderson, P. J., & Weissler, S. (1992). Recovery of coliforms from the hands of nurses and patients: Activities leading to contamination. *The Journal of Hospital Infection*, *21*(2), 85-93.
- Schneider, L. F., & Taylor, H. A. (1999). How do you get there from here? Mental representations of route descriptions. *Applied Cognitive Psychology* (Vol. 13, pp. 415-441): John Wiley & Sons Inc.
- Schneider, S. M., Prince-Paul, M., Allen, M. J., Silverman, P., & Talaba, D. (2004). Virtual reality as a distraction intervention for women receiving chemotherapy. *Oncology Nursing Forum*, *31*(1), 81-88.
- Schnelle, J. F., Ouslander, J. G., Simmons, S. F., Alessi, C. A., & Gravel, M. D. (1993). The nighttime environment, incontinence care, and sleep disruption in nursing homes. *Journal of the American Geriatrics Society*, *41*(9), 910-914.
- Sehulster, L., & Chinn, R. Y. (2003). Guidelines for environmental infection control in healthcare facilities. Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC). *MMWR Recommendation Report*, *52*(RR-10), 1-42.
- Shepley, M. M. (1995). The location of behavioral incidents in a children's psychiatric facility. *Children's Environments*, *12*(3), 352-361.
- Shepley, M. M. (2002). Predesign and postoccupancy analysis of staff behavior in a neonatal intensive care unit. *Children's Health Care*, *31*(3), 237-253.
- Shepley, M. M., & Davies, K. (2003). *Nursing Unit Configuration and Its Relationship to Noise and Nurse Walking Behavior: An AIDS/HIV Unit Case Study*. Retrieved 5/26/2004, from <http://www.aia.org/aah/journal/0401/article4.asp>

- Shepley, M. M., & Wilson, P. (1999). Designing for persons with AIDS: A post-occupancy study at the Bailey-Boushay House. *The Journal of Architectural & Planning Research*, 16(1), 17-32.
- Sherertz, R., J. Belani, A., Kramer, B. S., Elfenbein, G. J., Weiner, R. S., Sullivan, M. L., et al. (1987). Impact of air filtration on nosocomial *Aspergillus* infections: Unique risk of bone marrow transplant recipients. *American Journal of Medicine*, 83(4), 709-718.
- Sherertz, R. J., & Sullivan, M. L. (1985). An outbreak of infections with *Acinetobacter calcoaceticus* in burn patients: Contamination of patients' mattresses. *Journal of Infectious Diseases*, 151(2), 252-258.
- Shirani, K. Z., McManus, A. T., Vaughan, G. M., McManus, W. F., Pruitt, B. A., Jr., & Mason, A. D., Jr. (1986). Effects of environment on infection in burn patients. *Archives of Surgery*, 121(1), 31-36.
- Slevin, M., Farrington, N., Duffy, G., Daly, L., & Murphy, J. F. (2000). Altering the NICU and measuring infants' responses. *Acta Paediatrica*, 89(5), 577-581.
- Smedbold, H., Catrine, A., Unimed, S., Nilsen, A., Norback, D., & Hilt, B. (2002). Relationships between indoor environments and nasal inflammation in nursing personnel. *Archives of Environmental Health*, 57(2).
- Sommer, R., & Ross, H. (1958). Social interaction on a geriatrics ward. *The International Journal of Social Psychiatry*, 4(2), 128-133.
- Southwell, M. T., & Wistow, G. (1995). Sleep in hospitals at night: Are patients' needs being met? *Journal of Advanced Nursing*, 21(6), 1101-1109.
- Sturdavant, M. (1960). Intensive nursing service in circular and rectangular units. *Hospitals, JAHA*, 34(14), 46-48, 71-78.
- Swan, J. E., Richardson, L. D., & Hutton, J. D. (2003). Do appealing hospital rooms increase patient evaluations of physicians, nurses, and hospital services? *Health Care Management Review*, 28(3), 254.
- Terman, J. S., Terman, M., Lo, E.-S., & Cooper, T. B. (2001). Circadian time of morning light administration and therapeutic response in winter depression. *Archives of General Psychiatry*, 58(1), 69-75.
- Thompson, J. T., Meredith, J. W., & Molnar, J. A. (2002). The effect of burn nursing units on burn wound infections. *Journal of Burn Care Rehabilitation*, 23(4), 281-286.
- Topf, M. (1985). Noise-induced stress in hospital patients: Coping and nonauditory health outcomes. *Journal of Human Stress*, 11(3), 125-134.
- Topf, M., & Davis, J. E. (1993). Critical care unit noise and rapid eye movement (REM) sleep. *Heart Lung*, 22(3), 252-258.
- Topf, M., & Dillon, E. (1988). Noise-induced stress as a predictor of burnout in critical care nurses. *Heart Lung*, 17(5), 567-574.
- Topf, M., & Thompson, S. (2001). Interactive relationships between hospital patients' noise-induced stress and other stress with sleep. *Heart Lung*, 30(4), 237-243.
- Trites, D. K., Galbraith, F. D., Sturdavant, M., & Leckwart, J. F. (1970). Influence of nursing-unit design on the activities and subjective feelings of nursing personnel. *Environment & Behavior*, 2(3), 303-334.
- Tse, M. M.Y., Ng, J. K.F., Chung, J. W. Y., & Wong, T.K. S. (2002). The effect of visual stimuli on pain threshold and tolerance. *Journal of Clinical Nursing*, 11(4), 462-469.
- Ubel, P. A., Zell, M. M., & Miller, D. J. (1995). Elevator talk: Observational study of inappropriate comments in a public space. *American Journal of Medicine*, 99(2), 190-194.
- Uden, G. (1985). Inpatient accidents in hospitals. *Journal of the American Geriatrics Society*, 33(12), 833-841.
- Ulrich, R. S. (1984). View through a window may influence recovery from surgery. *Science*, 224(4647), 420-421.
- Ulrich, R. S. (1991). Effects of interior design on wellness: Theory and recent scientific research. *Journal of Health Care Interior Design*, 3(1), 97-109.
- Ulrich, R. S. (1999). Effects of gardens on health outcomes: Theory and research. In C. Cooper Marcus & M. Barnes (Eds.), *Healing gardens* (pp. 27-86). New York: Wiley.
- Ulrich, R. S., & Gilpin, L. (2003). Healing arts: Nutrition for the soul. In S. B. Frampton, L. Gilpin & P. Charmel (Eds.), *Putting patients first: Designing and practicing patient-centered care* (pp. 117-146). San Francisco: Jossey-Bass.

- Ulrich, R. S., Lawson, B., & Martinez, M. (2003). *Exploring the patient environment: An NHS estates workshop*. London: The Stationery Office.
- Ulrich, R. S., Simons, R. F., & Miles, M. A. (2003). Effects of environmental simulations and television on blood donor stress. *Journal of Architectural & Planning Research*, 20(1), 38-47.
- Van der Ploeg, H. M. (1988). Stressful medical events: A survey of patients' perceptions. In S. Maies, C. D. Spielberger, P. B. Defares & I. G. Sarason (Eds.), *Topics in Health Psychology* (pp. 193-203). New York: John Wiley.
- van Leeuwen, M., Bennett, L., West, S., Wiles, V., & Grasso, J. (2001). Patient falls from bed and the role of bedrails in the acute care setting. *Australian Journal of Advanced Nursing*, 19(2), 8-13.
- Van Someren, E. J. W., Kessler, A., Mirmiran, M., & Swaab, D. F. (1997). Indirect Bright Light Improves Circadian Rest-Activity Rhythm Disturbances in Demented Patients. *Biological Psychiatry*, 41(9), 955-963.
- Vassallo, M., Azeem, T., Pirwani, M. F., Sharma, J. C., & Allen, S. C. (2000). An epidemiological study of falls on integrated general medical wards. *International Journal of Clinical Practice*, 54(10), 654-657.
- Verderber, S. (1986). Dimensions of person-window transactions in the hospital environment. *Environment & Behavior*, 18(4), 450-466.
- Vernon, M. O., Trick, W. E., Welbel, S. F., Peterson, B. J., & Weinstein, R. A. (2003). Adherence with hand hygiene: Does number of sinks matter? *Infection Control and Hospital Epidemiology*, 24(3), 224-225.
- Volicer, B. J., Isenberg, M. A., & Burns, M. W. (1977). Medical-surgical differences in hospital stress factors. *Journal of Human Stress*, 3(2), 3-13.
- Walch, J. M., Rabin, B. S., Day, R., Williams, J. N., Choi, K., & Kang, J. D. (2004, In press). The effect of sunlight on post-operative analgesic medication usage: A prospective study of spinal surgery patients.
- Walder, B., Francioli, D., Meyer, J. J., Lancon, M., & Romand, J. A. (2000). Effects of guidelines implementation in a surgical intensive care unit to control nighttime light and noise levels. *Critical Care Medicine*, 28(7), 2242-2247.
- Wallace-Guy, G., Kripke, D., Jean-Louis, G., Langer, R., Elliott, J., & Tuunainen, A. (2002). Evening light exposure: Implications for sleep and depression. *Journal of the American Geriatrics Society*, 50(4), 738-739.
- Weisman, J. (1981). Evaluating architectural legibility: Wayfinding in the built environment. *Environment & Behavior*, 13(2), 189-204.
- Whall, A. L., Black, M. E., Groh, C. J., Yankou, D. J., Kupferschmid, B. J., & Foster, N. L. (1997). The effect of natural environments upon agitation and aggression in late stage dementia patients. *American Journal of Alzheimer's Disease and Other Dementias*, September-October, 216-220.
- Whitehouse, S., Varni, J. W., Seid, M., Cooper-Marcus, C., Ensberg, M. J., Jacobs, J. R., et al. (2001). Evaluating a children's hospital garden environment: Utilization and consumer satisfaction. *Journal of Environmental Psychology*, 21(3), 301-314.
- Williams, H. N., Singh, R., & Romberg, E. (2003). Surface contamination in the dental operator: A comparison over two decades. *Journal of the American Dental Association*, 134(3), 325-330;
- Wright, P., Hull, A. J., & Lickorish, A. (1993). Navigating in a hospital outpatients' department: The merits of maps and wallsigns. *Journal of Architectural and Planning research*, 10(1), 76-89.
- Yinnon, A. M., Ilan, Y., Tadmor, B., Altarescu, G., & Hershko, C. (1992). Quality of sleep in the medical department. *British Journal of Clinical Practitioners*, 46(2), 88-91.
- Zahr, L. K., & de Traversay, J. (1995). Premature infant responses to noise reduction by earmuffs: Effects on behavioral and physiologic measures. *Journal of Perinatology*, 15(6), 448-455.
- Zimring, C. (1990). *The costs of confusion: Non-monetary and monetary costs of the Emory University hospital wayfinding system*. Atlanta, GA: Georgia Institute of Technology.
- Zimring, C., & Templer, J. (1983-84). Wayfinding and orientation by the visually impaired. *Journal of Environmental Systems*, 13(4), 333-352.

## APPENDIX: Bibliography

### *Reduce Staff Stress, Fatigue; Increase Effectiveness in Delivering Care*

- Bookman, W. B. (1986). Space planning and staff turnover. *Certified Akers Laboratories*, 49(11), 22-23.
- Brennan, P. L., & Moos, R. H. (1990). Physical design, social climate, and staff turnover in skilled nursing facilities. *Journal of Long Term Care Administration*, 18(2), 22-27.
- Breslin, P. T., Gale, S. O., & Weitzner, W. M. (2003). Operational and design strategies for a reduced workforce. *Healthcare Financial Management*, 57(3), 62-65.
- Burgio, L., Engel, B., Hawkins, A., McCorick, K., & Scheve, A. (1990). A descriptive analysis of nursing staff behaviors in a teaching nursing home: Differences among NAs, LPNs and RNs. *Gerontologist*, 30(1), 107-112.
- Christensen, K. E. (1979). *An impact analysis framework for calculating the costs of staff disorientation in hospitals*. Los Angeles: School of Architecture and Urban Planning, University of California.
- Dennis, K. E., Sweeney, P. M., Macdonald, L. P., & Morse, N. A. (1993). Point of care technology: Impact on people and paperwork. *Nursing Economics*, 11(4), 229-237, 248.
- Garg, A., & Owen, B. (1992). Reducing back stress to nursing personnel: An ergonomic intervention in a nursing home. *Ergonomics*, 35(11), 1353-1375.
- Hibbard, F. J., Bosso, J. A., Sward, L. W., & Baum, S. (1981). Delivery time in a decentralized pharmacy system without satellites. *American Journal of Hospital Pharmacy*, 38(5), 690-692.
- Janssen, P. A., Harris, S. J., Soolsma, J., Klein, M. C., & Seymour, L. C. (2001). Single room maternity care: The nursing response. *Birth*, 28(3), 173-179.
- Jiang, S., Huang, L., Chen, X., Wang, J., Wu, W., Yin, S., et al. (2003). Ventilation of wards and nosocomial outbreak of severe acute respiratory syndrome among healthcare workers. *Chinese Medical Journal*, 116(9), 1293-1297.
- Joint Commission on Accreditation of Healthcare Organizations (2002). *Health care at the crossroads: Strategies for addressing the evolving nursing crisis*. Oakbrook Terrace, IL: Joint Commission on Accreditation of Healthcare Organizations.
- Jones, R. G. (1988). Experimental study to evaluate nursing staff morale in a high stimulation geriatric psychiatry setting. *Journal of Advanced Nursing*, 13(3), 352-357.
- Klimowski, L. L., Rotstein, C., & Cummings, K. M. (1989). Incidence of nosocomial aspergillosis in patients with leukemia over a twenty-year period. *Infection Control and Hospital Epidemiology*, 10(7), 299-305.
- Kromhout, H., Hoek, F., Uitterhoeve, R., Huijbers, R., Overmars, R. F., Anzion, R., et al. (2000). Postulating a dermal pathway for exposure to anti-neoplastic drugs among hospital workers: Applying a conceptual model to the results of three workplace surveys. *Annals of Occupational Hygiene*, 44(7), 551-560.
- Lin, A. C., Barker, K. N., Hassall, T. H., & Gallelli, J. F. (1988). Effects of simulated facility-design changes on outpatient pharmacy efficiency. *American Journal of Hospital Pharmacy*, 45(1), 116-121.
- Lin, A. C., Jang, R., Sedani, D., Thomas, S., Barker, K. N., & Flynn, E. A. (1996). Re-engineering a pharmacy work system and layout to facilitate patient counseling. *American Journal of Health-System Pharmacy*, 53(13), 1558-1564.
- Marr, P. B., Duthie, E., Glassman, K. S., Janovas, D. M., Kelly, J. B., Graham, E., et al. (1993). Bedside terminals and quality of nursing documentation. *Computers in Nursing*, 11(4),

- 176-182.
- Pierce, R. A., 2nd, Rogers, E. M., Sharp, M. H., & Musulin, M. (1990). Outpatient pharmacy redesign to improve work flow, waiting time, and patient satisfaction. *American Journal of Hospital Pharmacy*, 47(2), 351-356.
- Reynolds, D. M., Johnson, M. H., & Longe, R. L. (1978). Medication delivery time requirements in centralized and decentralized unit dose drug distribution systems. *American Journal of Hospital Pharmacy*, 35(8), 941-943.
- Shepley, M. M., & Davies, K. (2003). *Nursing unit configuration and its relationship to noise and nurse walking behavior: An AIDS/HIV unit case study*. Retrieved May 26, 2004, from <http://www.aia.org/aah/journal/0401/article4.asp>.
- Smedbold, H. T., Ahlen, C., Unimed, S., Nilsen, A. M., Norbaeck, D., & Hilt, B. (2002). Relationships between indoor environments and nasal inflammation in nursing personnel. *Archives of Environmental Health*, 57(2), 155-161.
- Stefanchik, M. F., & Cohen, P. I. (1989). Point-of-care faces cost challenges. *Computers in Healthcare*, 10(12), 38-39, 41.
- Summer, G., Lirk, P., Hoerauf, K., Riccabona, U., Bodrogi, F., Raifer, H., et al. (2003). Sevoflurane in exhaled air of operating room personnel. *Anesthesia and Analgesia*, 97(4), 1070.
- Sutton, P. M., Mossman, M., Reinisch, F., & Harrison, R. J. (2002). A determination of healthcare worker exposure to Mycobacterium tuberculosis in two Californian hospitals. *Journal of Hospital Infection*, 51(4), 281-287.
- Trites, D. K., Galbraith, F. D., Sturdavant, M., & Leckwart, J. F. (1970). Influence of nursing-unit design on the activities and subjective feelings of nursing personnel. *Environment & Behavior*, 2(3), 303-334.
- Tyson, G. A., Lambert, G., & Beattie, L. (2002). The impact of ward design on the behaviour, occupational satisfaction and well-being of psychiatric nurses. *International Journal of Mental Health Nursing*, 11(2), 94-102.
- Walker, S. B. (1998). Neonatal nurses' views on the barriers to parenting in the intensive care nursery—a national study. *Australian Critical Care*, 11(3), 86-91.
- Willson, D. (1994). Survey of nurse perceptions regarding the utilization of bedside computers. *Proceedings of Annual Symposium on Computer Application in Medical Care*, 18, 553-557.

### ***Improve Patient Safety***

- Abzug, M. J., Gardner, S., Glode, M. P., Cymanski, M., Roe, M. H., & Odom, L. F. (1992). Heliport-associated nosocomial mucormycoses. *Infection Control and Hospital Epidemiology*, 13(6), 325-326.
- Adeniran, A., Shakespeare, P., Patrick, S., Fletcher, A. J., & Rossi, L. A. (1995). Influence of a changed care environment on bacterial colonization of burn wounds. *Burns*, 21(7), 521-525.
- Albert, R. K., & Condie, F. (1981). Hand-washing patterns in medical intensive-care units. *New England Journal of Medicine*, 304(24), 1465-1466.
- Alberti, C., Bouakline, A., Ribaud, P., Lacroix, C., Rousselot, P., Leblanc, T., et al. (2001). Relationship between environmental fungal contamination and the incidence of invasive aspergillosis in haematology patients. *Journal of Hospital Infection*, 48(3), 198-206.
- Alcee, D. A. (2000). The experience of a community hospital in quantifying and reducing patient falls. *Journal of Nursing Care Quality*, 14(3), 43-54.
- Allander, T., Gruber, A., Naghavi, M., Beyene, A., Soderstrom, T., Bjorkholm, M., et al. (1995).

- Frequent patient-to-patient transmission of hepatitis C virus in a haematology ward. *Lancet*, 345(8950), 603-607.
- American Health Consultants (1995). Aspergillosis: A deadly dust may be in the wind during renovations. *Hospital Infection Control*, 22(10), 125-126.
- Andersen, B. M., & Rasch, M. (2000). Hospital-acquired infections in Norwegian long-term-care institutions. A three-year survey of hospital-acquired infections and antibiotic treatment in nursing/residential homes, including 4500 residents in Oslo. *Journal of Hospital Infection*, 46(4), 288-296.
- Anderson, J. D., Bonner, M., Scheifele, D. W., & Schneider, B. C. (1985). Lack of nosocomial spread of Varicella in a pediatric hospital with negative pressure ventilated patient rooms. *Infection Control*, 6(3), 120-121.
- Anderson, K., Morris, G., Kennedy, H., Croall, J., Michie, J., Richardson, M., et al. (1996). Aspergillosis in immunocompromised paediatric patients: Associations with building hygiene, design and indoor air. *Thorax*, 51(3), 256-261.
- Anderson, R. L., Mackel, D. C., Stoler, B. S., & Mallison, G. F. (1982). Carpeting in hospitals: An epidemiological evaluation. *Journal of Clinical Microbiology*, 15(3), 408-415.
- Archibald, N. (2000). Preventing serious injuries from falls. *Provider*, 26(Suppl. 9), 8-9, 16.
- Archibald, L. K., Manning, M. L., Bell, L. M., Banerjee, S., & Jarvis, W. R. (1997). Patient density, nurse-to-patient ratio and nosocomial infection risk in a pediatric cardiac intensive care unit. *Pediatric Infectious Disease Journal*, 16(11), 1045-1048.
- Arlet, G., Gluckman, E., Gerber, F., Perol, Y., & Hirsch, A. (1989). Measurement of bacterial and fungal air counts in two bone marrow transplant units. *Journal of Hospital Infection*, 13(1), 63-69.
- Armstrong, D. (1984). Protected environments are discomfoting and expensive and do not offer meaningful protection. *American Journal of Medicine*, 76(4), 685-689.
- Ash, K. L., MacLeod, P., & Clark, L. (1998). A case control study of falls in the hospital setting. *Journal of Gerontological Nursing*, 24(12), 7-15.
- Austin, D. J., Bonten, M. J., Weinstein, R. A., Slaughter, S., & Anderson, R. M. (1999). Vancomycin-resistant enterococci in intensive-care hospital settings: Transmission dynamics, persistence, and the impact of infection control programs. *Proceedings of the National Academy of Sciences of the United States of America*, 96(12), 6908-6913.
- Aygun, G., Demirkiran, O., Utku, T., Mete, B., Urkmez, S., Yilmaz, M., et al. (2002). Environmental contamination during a carbapenem-resistant *Acinetobacter baumannii* outbreak in an intensive care unit. *Journal of Hospital Infection*, 52(4), 259-262.
- Babb, J. R., Lynam, P., & Ayliffe, G. A. (1995). Risk of airborne transmission in an operating theatre containing four ultraclean air units. *Journal of Hospital Infection*, 31(3), 159-168.
- Bakarich, A., McMillan, V., & Prosser, R. (1997). The effect of a nursing intervention on the incidence of older patient falls. *Australian Journal of Advanced Nursing*, 15(1), 26-31.
- Barber, N. D., Blackett, A., & Batty, R. (1993). Does a high workload decrease ward pharmacists' clinical monitoring? *International Journal of Pharmacy Practice*, 2, 152-155.
- Barlas, D., Sama, A. E., Ward, M. F., & Lesser, M. L. (2001). Comparison of the auditory and visual privacy of emergency department treatment areas with curtains versus those with solid walls. *Annals of Emergency Medicine*, 38(2), 135-139.
- Barnes, R. A., & Rogers, T. R. (1989). Control of an outbreak of nosocomial aspergillosis by laminar air-flow isolation. *Journal of Hospital Infection*, 14(2), 89-94.
- Bauer, T. M., Ofner, E., Just, H. M., Just, H., & Daschner, F. D. (1990). An epidemiological study assessing the relative importance of airborne and direct contact transmission of microorganisms in a medical intensive care unit. *Journal of Hospital Infection*, 15(4), 301-309.
- Bayat, A., Shaaban, H., Dodgson, A., & Dunn, K. W. (2003). Implications for burns unit design

- following outbreak of multi-resistant *Acinetobacter* infection in ICU and burns unit. *Burns*, 29(4), 303-306.
- Becker, C., Kron, M., Lindemann, U., Sturm, E., Eichner, B., Walter-Jung, B., et al. (2003). Effectiveness of a multifaceted intervention on falls in nursing home residents. *Journal of the American Geriatrics Society*, 51(3), 306-313.
- Ben-Abraham, R., Keller, N., Szold, O., Vardi, A., Weinberg, M., Barzilay, Z., et al. (2002). Do isolation rooms reduce the rate of nosocomial infections in the pediatric intensive care unit? *Journal of Critical Care*, 17(3), 176-180.
- Bertrand, X., Bailly, P., Blasco, G., Balvay, P., Boillot, A., & Talon, D. (2000). Large outbreak in a surgical intensive care unit of colonization or infection with *Pseudomonas aeruginosa* that overexpressed an active efflux pump. *Clinical Infectious Diseases*, 31, E9-E14.
- Beyer, D. J., & Belsito, D. V. (2000). Fungal contamination of outpatient examination rooms: Is your office safe? *Dermatology Nursing*, 12(1), 51-53.
- Bischoff, W. E., Reynolds, T. M., Sessler, C. N., Edmond, M. B., & Wenzel, R. P. (2000). Handwashing compliance by health care workers: The impact of introducing an accessible, alcohol-based hand antiseptic. *Archives of Internal Medicine*, 160(7), 1017-1021.
- Bittner, M. J., & Rich, E. C. (1998). Surveillance of handwashing episodes in adult intensive-care units by measuring an index of soap and paper towel consumption. *Clinical Performance and Quality Health Care*, 6(4), 179-182.
- Bjorn, E., & Nielsen, P. V. (2002). Dispersal of exhaled air and personal exposure in displacement ventilated rooms. *Indoor Air*, 12(3), 147-164.
- Bodey, G. P., & Johnston, D. (1971). Microbiological evaluation of protected environments during patient occupancy. *Applied Microbiology*, 22(5), 828-836.
- Booker, J. M., & Roseman, C. (1995). A seasonal pattern of hospital medication errors in Alaska. *Psychiatry Research*, 57(3), 251-257.
- Borg, M. A. (2003). Bed occupancy and overcrowding as determinant factors in the incidence of MRSA infections within general ward settings. *Journal of Hospital Infection*, 54(4), 316-318.
- Boyce, J. M., Potter-Bynoe, G., Chenevert, C., & King, T. (1997). Environmental contamination due to methicillin-resistant *Staphylococcus aureus*: Possible infection control implications. *Infection Control and Hospital Epidemiology*, 18(9), 622-627.
- Brandis, S. (1999). A collaborative occupational therapy and nursing approach to falls prevention in hospital inpatients. *Journal of Quality in Clinical Practice*, 19(4), 215-221.
- Breiman, R., Cozen, W., & Fields, B. (1990). Role of air sampling in investigation of an outbreak of Legionnaires' disease associated with exposure to aerosols from an evaporative condenser. *Journal of Infectious Diseases*, 161(6), 1257-1261.
- Brown, I. W., Jr., Moor, G. F., Hummel, B. W., Marshall, W. G., Jr., & Collins, J. P. (1996). Toward further reducing wound infections in cardiac operations. *Annals of Thoracic Surgery*, 62(6), 1783-1789.
- Bruce, B. B., Blass, M. A., Blumberg, H. M., Lennox, J. L., del Rio, C., & Horsburgh, C. R., Jr. (2000). Risk of *Cryptosporidium parvum* transmission between hospital roommates. *Clinical Infectious Diseases*, 31(4), 947-950.
- Buchanan, T. L., Barker, K. N., Gibson, J. T., Jiang, B. C., & Pearson, R. E. (1991). Illumination and errors in dispensing. *American Journal of Hospital Pharmacy*, 48(10), 2137-2145.
- Bures, S., Fishbain, J. T., Uyehara, C. F., Parker, J. M., & Berg, B. W. (2000). Computer keyboards and faucet handles as reservoirs of nosocomial pathogens in the intensive care unit. *American Journal of Infection Control*, 28(6), 465-471.
- Burke, J. F., Quinby, W. C., Bondoc, C. C., Sheehy, E. M., & Moreno, H. C. (1977). The contribution of a bacterially isolated environment to the prevention of infection in



- seriously burned patients. *Annals of Surgery*, 186(3), 377-387.
- Capezuti, E., Maislin, G., Strumpf, N., & Evans, L. K. (2002). Side rail use and bed-related fall outcomes among nursing home residents. *Journal of the American Geriatrics Society*, 50(1), 90-96.
- Carducci, A., Verani, M., Casini, B., Giuntini, A., Mazzoni, F., Rovini, E., et al. (2002). Detection and potential indicators of the presence of hepatitis C virus on surfaces in hospital settings. *Letters in Applied Microbiology*, 34(3), 189-193.
- Carter, C. D., & Barr, B. A. (1997). Infection control issues in construction and renovation. *Infection Control and Hospital Epidemiology*, 18(8), 587-596.
- Chang, V. T., & Nelson, K. (2000). The role of physical proximity in nosocomial diarrhea. *Clinical Infectious Diseases*, 31(3), 717-722.
- Chen, Q., Jiang, Z., & Moser, A. (1992). Control of airborne particle concentration and draught risk in an operating room. *Indoor Air*, 2(3), 154-167.
- Cheng, V. C., Lo, W. K., Woo, P. C., Chan, S. B., Cheng, S. W., Ho, M., et al. (2001). Polymicrobial outbreak of intermittent peritoneal dialysis peritonitis during external wall renovation at a dialysis center. *Peritoneal Dialysis International*, 21(3), 296-301.
- Cohen, B., Saiman, L., Cimiotti, J., & Larson, E. (2003). Factors associated with hand hygiene practices in two neonatal intensive care units. *Pediatric Infectious Disease Journal*, 22(6), 494-499.
- Cohen, H. A., Kitai, E., Levy, I., & Ben-Amitai, D. (2002). Handwashing patterns in two dermatology clinics. *Dermatology*, 205(4), 358-361.
- Conly, J. M., Hill, S., Ross, J., Lertzman, J., & Louie, T. J. (1989). Handwashing practices in an intensive care unit: The effects of an educational program and its relationship to infection rates. *American Journal of Infection Control*, 17(6), 330-339.
- Connell, B. R. (1996). Role of the environment in falls prevention. *Clinics in Geriatric Medicine*, 12(4), 859-880.
- Cook, R. I., Render, M., & Woods, D. D. (2000). Gaps in the continuity of care and progress on patient safety. *British Medical Journal (Clinical Research Ed.)*, 320(7), 791-794.
- Cornet, M., Levy, V., Fleury, L., Lortholary, J., Barquins, S., Coureul, M. H., et al. (1999). Efficacy of prevention by high-efficiency particulate air filtration or laminar airflow against *Aspergillus* airborne contamination during hospital renovation. *Infection Control and Hospital Epidemiology*, 20(7), 508-513.
- Cotterill, S., Evans, R., & Fraise, A. P. (1996). An unusual source for an outbreak of methicillin-resistant *Staphylococcus aureus* on an intensive therapy unit. *Journal of Hospital Infection*, 32(3), 207-216.
- Davidson, A. I., Smylie, H. G., Macdonald, A., & Smith, G. (1971). Ward design in relation to postoperative wound infection. *British Medical Journal*, 1(740), 72-75.
- Dean, B., Schachter, M., Vincent, C., & Barber, N. (2002). Causes of prescribing errors in hospital inpatients: A prospective study. *Lancet*, 359(9315), 1373-1378.
- Dekker, A. W., Verdonck, L. F., & Rozenberg-Arska, M. (1994). Infection prevention in autologous bone marrow transplantation and the role of protective isolation. *Bone Marrow Transplantation*, 14(1), 89-93.
- Demling, R., Perea, A., Maly, J., Moylan, J., Jarett, F., & Balish, E. (1978). The use of a laminar airflow isolation system for the treatment of major burns. *American Journal of Surgery*, 136(3), 375-378.
- deSilva, M. I., & Rissing, J. P. (1984). Postoperative wound infections following cardiac surgery: Significance of contaminated cases performed in the preceding 48 hours. *Infection Control*, 5(8), 371-377.
- Dettenkofer, M., Scherrer, M., Hoch, V., Glaser, H., Schwarzer, G., Zentner, J., et al. (2003). Shutting down operating theater ventilation when the theater is not in use: Infection

- control and environmental aspects. *Infection Control and Hospital Epidemiology*, 24(8), 596-600.
- Devine, J., Cooke, R. P., & Wright, E. P. (2001). Is methicillin-resistant *Staphylococcus aureus* (MRSA) contamination of ward-based computer terminals a surrogate marker for nosocomial MRSA transmission and handwashing compliance? *Journal of Hospital Infection*, 48(1), 72-75.
- Dharan, S., & Pittet, D. (2002). Environmental controls in operating theatres. *Journal of Hospital Infection*, 51(2), 79-84.
- Doebbeling, B. N., Stanley, G. L., Sheetz, C. T., Pfaller, M. A., Houston, A. K., Annis, L., et al. (1992). Comparative efficacy of alternative hand-washing agents in reducing nosocomial infections in intensive care units. *New England Journal of Medicine*, 327(2), 88-93.
- Donald, I. P., Pitt, K., Armstrong, E., & Shuttleworth, H. (2000). Preventing falls on an elderly care rehabilitation ward. *Clinical Rehabilitation*, 14(2), 178-185.
- Donowitz, L. G. (1987). Handwashing technique in a pediatric intensive care unit. *American Journal of Diseases of Children*, 141(6), 683-685.
- Dorsey, S. T., Cydulka, R. K., & Emerman, C. L. (1996). Is handwashing teachable? Failure to improve handwashing behavior in an urban emergency department. *Academic Emergency Medicine*, 3(4), 360-365.
- Dubbert, P. M., Dolce, J., Richter, W., Miller, M., & Chapman, S. W. (1990). Increasing ICU staff handwashing: Effects of education and group feedback. *Infection Control and Hospital Epidemiology*, 11(4), 191-193.
- Enser, J., & MacInnes, D. (1999). The relationship between building design and escapes from secure units. *Journal of the Royal Society of Health*, 119(3), 170-174.
- Farquharson, C., & Baguley, K. (2003). Responding to the severe acute respiratory syndrome (SARS) outbreak: Lessons learned in a Toronto emergency department. *Journal of Emergency Nursing*, 29(3), 222-228.
- Flynn, E. A., Barker, K. N., Gibson, J. T., Pearson, R. E., Berger, B. A., & Smith, L. A. (1999). Impact of interruptions and distractions on dispensing errors in an ambulatory care pharmacy. *American Journal of Health-System Pharmacy*, 56(13), 1319-1325.
- Friberg, B., Friberg, S., & Burman, L. G. (1999). Correlation between surface and air counts of particles carrying aerobic bacteria in operating rooms with turbulent ventilation: An experimental study. *Journal of Hospital Infection*, 42(1), 61-68.
- Friberg, S., Ardnor, B., & Lundholm, R. (2003). The addition of a mobile ultra-clean exponential laminar airflow screen to conventional operating room ventilation reduces bacterial contamination to operating box levels. *Journal of Hospital Infection*, 55(2), 92-97.
- Fridkin, S. K., Kremer, F. B., Bland, L. A., Padhye, A., McNeil, M. M., & Jarvis, W. R. (1996). *Acromonium kiliense* endophthalmitis that occurred after cataract extraction in an ambulatory surgical center and was traced to an environmental reservoir. *Clinical Infectious Diseases*, 22(2), 222-227.
- Gardner, P. S., Court, S. D., Brocklebank, J. T., Downham, M. A., & Weightman, D. (1973). Virus cross-infection in paediatric wards. *British Medical Journal*, 2(5866), 571-575.
- Garner, J. S. (1996). Guideline for isolation precautions in hospitals. *Infection Control and Hospital Epidemiology*, 17(1), 53-80.
- Gerson, S. L., Parker, P., Jacobs, M. R., Creger, R., & Lazarus, H. M. (1994). Aspergillosis due to carpet contamination. *Infection Control and Hospital Epidemiology*, 15(1), 221-223.
- Goldmann, D. A., Durbin, W. A., Jr., & Freeman, J. (1981). Nosocomial infections in a neonatal intensive care unit. *Journal of Infectious Diseases*, 144(5), 449-459.
- Goodley, J. M., Clayton, Y. M., & Hay, R. J. (1994). Environmental sampling for aspergilli during building construction on a hospital site. *Journal of Hospital Infection*, 26(1), 27-35.

- Graham, M. (1990). Frequency and duration of handwashing in an intensive care unit. *American Journal of Infection Control*, 18(2), 77-81.
- Greenfield, E., & McManus, A. T. (1997). Infectious complications: Prevention and strategies for their control. *Nursing Clinics of North America*, 32(2), 297-309.
- Guernsey, B. G., Ingram, N. B., Hokanson, J. A., Doutre, W. H., Bryant, S. G., Blair, C. W., et al. (1983). Pharmacists' dispensing accuracy in a high-volume outpatient pharmacy service: Focus on risk management. *Drug Intelligence & Clinical Pharmacy*, 17(10), 742-746.
- Hahn, T., Cummings, K. M., Michalek, A. M., Lipman, B. J., Segal, B. H., & McCarthy, P. L., Jr. (2002). Efficacy of high-efficiency particulate air filtration in preventing aspergillosis in immunocompromised patients with hematologic malignancies. *Infection Control and Hospital Epidemiology*, 23(9), 525-531.
- Hamrick, W. B., & Reilly, L. (1992). A comparison of infection rates in a newborn intensive care unit before and after adoption of open visitation. *Neonatal Network* 11(1), 15-18.
- Hanger, H. C., Ball, M. C., & Wood, L. A. (1999). An analysis of falls in the hospital: Can we do without bedrails? *Journal of the American Geriatrics Society*, 47(5), 529-531.
- Hannan, M. M., Azadian, B. S., Gazzard, B. G., Hawkins, D. A., & Hoffman, P. N. (2000). Hospital infection control in an era of HIV infection and multi-drug resistant tuberculosis. *Journal of Hospital Infection*, 44(1), 5-11.
- Harbarth, S., Sudre, P., Dharan, S., Cadenas, M., & Pittet, D. (1999). Outbreak of *Enterobacter cloacae* related to understaffing, overcrowding, and poor hygiene practices. *Infection Control and Hospital Epidemiology*, 20(9), 598-603.
- Harvey, M. A. (1998). Critical-care-unit bedside design and furnishing: Impact on nosocomial infections. *Infection Control and Hospital Epidemiology*, 19(8), 597-601.
- Hendrich, A., Fay, J., & Sorrells, A. (2002, September). Courage to heal: Comprehensive cardiac critical care. *Healthcare Design*, 11-13.
- Hendrich, A. (2003, October). *Optimizing physical space for improved outcomes: Satisfaction and the bottom line*. Paper presented at "Optimizing the Physical Space for Improved Outcomes, Satisfaction, and the Bottom Line," minicourse sponsored by the Institute for Healthcare Improvement and the Center for Health Design, Atlanta, GA.
- Hendrich, A. L., Bender, P. S., & Nyhuis, A. (2003). Validation of the Hendrich II Fall Risk Model: A large concurrent case/control study of hospitalized patients. *Applied Nursing Research*, 16(1), 9-21.
- Herrero, I. A., Issa, N. C., & Patel, R. (2002). Nosocomial spread of linezolid-resistant, vancomycin-resistant *Enterococcus faecium*. *New England Journal of Medicine*, 346(11), 867-869.
- Hofmann, M. T., Bankes, P. F., Javed, A., & Selhat, M. (2003). Decreasing the incidence of falls in the nursing home in a cost-conscious environment: A pilot study. *Journal of the American Medical Directors Association*, 4(2), 95-97.
- Hopkins, C. C., Weber, D. J., & Rubin, R. H. (1989). Invasive aspergillus infection: Possible non-ward common source within the hospital environment. *Journal of Hospital Infection*, 13(1), 19-25.
- Huebner, J., Frank, U., & Kappstein, I. (1989). Influence of architectural design on nosocomial infections in intensive care units. *Intensive Care Medicine*, 15(3), 179-183.
- Humphreys, H., Johnson, E. M., Warnock, D. W., Willatts, S. M., Winter, R. J., & Speller, D. C. (1991). An outbreak of aspergillosis in a general ITU. *Journal of Hospital Infection*, 18(3), 167-177.
- Institute of Medicine. (2004). *Keeping patients safe: Transforming the work environment of nurses*. Washington, DC: National Academy Press.
- Iwen, P. C., Davis, J. C., Reed, E. C., Winfield, B. A., & Hinrichs, S. H. (1994). Airborne fungal spore monitoring in a protective environment during hospital construction, and

- correlation with an outbreak of invasive aspergillosis. *Infection Control and Hospital Epidemiology*, 15(5), 303-306.
- Jack, C. I., Smith, T., Neoh, C., Lye, M., & McGalliard, J. N. (1995). Prevalence of low vision in elderly patients admitted to an acute geriatric unit in Liverpool: Elderly people who fall are more likely to have low vision. *Gerontology*, 41(5), 280-285.
- Jensen, J., Lundin-Olsson, L., Nyberg, L., & Gustafson, Y. (2002). Falls among frail older people in residential care. *Scandinavian Journal of Public Health*, 30(1), 54-61.
- Jensen, J., Nyberg, L., Gustafson, Y., & Lundin-Olsson, L. (2003). Fall and injury prevention in residential care—effects in residents with higher and lower levels of cognition. *Journal of the American Geriatrics Society*, 51(5), 627-635.
- Jernigan, J. A., Titus, M. G., Groschel, D. H., Getchell-White, S., & Farr, B. M. (1996). Effectiveness of contact isolation during a hospital outbreak of methicillin-resistant *Staphylococcus aureus*. *American Journal of Epidemiology*, 143(5), 496-504.
- Kaplan, L. M., & McGuckin, M. (1986). Increasing handwashing compliance with more accessible sinks. *Infection Control*, 7(8), 408-410.
- Kates, S. G., McGinley, K. J., Larson, E. L., & Leyden, J. J. (1991). Indigenous multiresistant bacteria from flowers in hospital and nonhospital environments. *American Journal of Infection Control*, 19(3), 156-161.
- Kim, M. H., Mindorff, C., Patrick, M. L., Gold, R., & Ford-Jones, E. L. (1987). Isolation usage in a pediatric hospital. *Infection Control*, 8(5), 195-199.
- Kistner, U. A., Keith, M. R., Sergeant, K. A., & Hokanson, J. A. (1994). Accuracy of dispensing in a high-volume, hospital-based outpatient pharmacy. *American Journal of Hospital Pharmacy*, 51(22), 2793-2797.
- Kumari, D. N., Haji, T. C., Keer, V., Hawkey, P. M., Duncanson, V., & Flower, E. (1998). Ventilation grilles as a potential source of methicillin-resistant *Staphylococcus aureus* causing an outbreak in an orthopaedic ward at a district general hospital. *Journal of Hospital Infection*, 39(2), 127-133.
- Lai, K. K. (2001). A cluster of invasive aspergillosis in a bone marrow transplant unit related to construction and the utility of air sampling. *American Journal of Infection Control*, 29(5), 333-337.
- Langley, J. M., Hanakowski, M., & Bortolussi, R. (1994). Demand for isolation beds in a pediatric hospital. *American Journal of Infection Control*, 22(4), 207-211.
- Lankford, M. G., Zembower, T. R., Trick, W. E., Hacek, D. M., Noskin, G. A., & Peterson, L. R. (2003). Influence of role models and hospital design on hand hygiene of healthcare workers. *Emerging Infectious Diseases*, 9(2), 217-223.
- LaPointe, N. M., & Jollis, J. G. (2003). Medication errors in hospitalized cardiovascular patients. *Archives of Internal Medicine*, 163(12), 1461-1466.
- Larson, E. (1988). A causal link between handwashing and risk of infection? Examination of the evidence. *Infection Control*, 9(1), 28-36.
- Larson, E., & Kretzer, E. K. (1995). Compliance with handwashing and barrier precautions. *Journal of Hospital Infection*, 30 (Suppl.), 88-106.
- Larson, E., McGeer, A., Quraishi, Z. A., Krenzischek, D., Parsons, B. J., Holdford, J., et al. (1991). Effect of an automated sink on handwashing practices and attitudes in high-risk units. *Infection Control and Hospital Epidemiology*, 12(7), 422-428.
- Larson, E. L., Bryan, J. L., Adler, L. M., & Blane, C. (1997). A multifaceted approach to changing handwashing behavior. *American Journal of Infection Control*, 25(1), 3-10.
- Larson, E. L., McGinley, K. J., Foglia, A., Leyden, J. J., Boland, N., Larson, J., et al. (1992). Handwashing practices and resistance and density of bacterial hand flora on two pediatric units in Lima, Peru. *American Journal of Infection Control*, 20(2), 65-72.
- Laurel, V. L., Meier, P. A., Astorga, A., Dolan, D., Brockett, R., & Rinaldi, M. G. (1999).

- Pseudoepidemic of *Aspergillus niger* infections traced to specimen contamination in the microbiology laboratory. *Journal of Clinical Microbiology*, 37(5), 1612-1616.
- Layton, M. C., Perez, M., Heald, P., & Patterson, J. E. (1993). An outbreak of mupirocin-resistant *Staphylococcus aureus* on a dermatology ward associated with an environmental reservoir. *Infection Control and Hospital Epidemiology*, 14(7), 369-375.
- Leclair, J. M., Freeman, J., Sullivan, B. F., Crowley, C. M., & Goldmann, D. A. (1987). Prevention of nosocomial respiratory syncytial virus infections through compliance with glove and gown isolation precautions. *New England Journal of Medicine*, 317(6), 329-334.
- Lee, M. L., Chen, C. J., Su, I., Chen, K. T., Yeh, C. C., King, C. C., Chang, H. I., et al. (2003). Use of quarantine to prevent transmission of severe acute respiratory syndrome—Taiwan, 2003. *MMWR*, 52(29), 680-683.
- Lesar, T. S., Briceland, L. L., Delcours, K., Parmalee, J. C., Masta-Gornic, V., & Pohl, H. (1990). Medication prescribing errors in a teaching hospital. *Journal of the American Medical Association*, 263(17), 2329-2334.
- Levine, A., Siegel, S., Schreiber, A., Hauser, J., Preisler, H., Goldstein, I. M., et al. (1973). Protected environments and prophylactic antibiotics. *New England Journal of Medicine*, 288(10), 477-483.
- Li, C.-S., & Hou, P.-A. (2003). Bioaerosol characteristics in hospital clean rooms. *The Science of the Total Environment*, 305(3), 169-176.
- Lidwell, O. M. (1976, October). Clean air, less infection. *Hospital Engineering*, 9-17.
- Lidwell, O. M., Lowbury, E. J., Whyte, W., Blowers, R., Stanley, S. J., & Lowe, D. (1984). Infection and sepsis after operations for total hip or knee-joint replacement: Influence of ultraclean air, prophylactic antibiotics and other factors. *Journal of Hygiene*, 93(3), 505-529.
- Loeb, M., Wilcox, L., Thornley, D., Gun-Munro, J., & Richardson, H. (1995). *Bacillus* species pseudobacteremia following hospital construction. *Canadian Journal of Infection Control*, 10(2), 37-40.
- Loo, V. G., Bertrand, C., Dixon, C., Vitye, D., DeSalis, B., McLean, A. P., et al. (1996). Control of construction-associated nosocomial aspergillosis in an antiquated hematology unit. *Infection Control and Hospital Epidemiology*, 17(6), 360-364.
- Lord, S. R., Clark, R. D., & Webster, I. W. (1991). Visual acuity and contrast sensitivity in relation to falls in an elderly population. *Age and Ageing*, 20(3), 175-181.
- Lowery, K., Buri, H., & Ballard, C. (2000). What is the prevalence of environmental hazards in the homes of dementia sufferers and are they associated with falls? *International Journal of Geriatric Psychiatry*, 15(10), 883-886.
- Lund, S., Jackson, J., Leggett, J., Hales, L., Dworkin, R., & Gilbert, D. (1994). Reality of glove use and handwashing in a community hospital. *American Journal of Infection Control*, 22(6), 352-357.
- Lutz, B. D., Jiankang, J., Rinaldi, M. G., Wickes, B. L., & Huycke, M. M. (2003). Outbreak of invasive *Aspergillus* infection in surgical patients, associated with a contaminated air-handling system. *Clinical Infectious Diseases*, 37(6), 786-793.
- Mahieu, L. M., De Dooy, J. J., Van Laer, F. A., Jansens, H., & Ieven, M. M. (2000). A prospective study on factors influencing aspergillus spore load in the air during renovation works in a neonatal intensive care unit. *Journal of Hospital Infection*, 45(3), 191-197.
- Maki, D. G. (1989). Risk factors for nosocomial infection in intensive care. 'Devices vs nature' and goals for the next decade. *Archives of Internal Medicine*, 149(1), 30-35.
- Maki, D. G., Alvarado, C. J., Hassemer, C. A., & Zilz, M. A. (1982). Relation of the inanimate hospital environment to endemic nosocomial infection. *New England Journal of*

- Medicine*, 307(25), 1562-1566.
- Malamou-Ladas, H., O'Farrell, S., Nash, J. Q., & Tabaqchali, S. (1983). Isolation of *Clostridium difficile* from patients and the environment of hospital wards. *Journal of Clinical Pathology*, 36(1), 88-92.
- Malik, R. K., Montecalvo, M. A., Reale, M. R., Li, K., Maw, M., Munoz, J. L., et al. (1999). Epidemiology and control of vancomycin-resistant enterococci in a regional neonatal intensive care unit. *Pediatric Infectious Disease Journal*, 18(4), 352-356.
- Marier, R. L. (1996). The design of isolation rooms. *Infection Control and Hospital Epidemiology*, 17(1), 3-4.
- McDonald, L. C., Walker, M., Carson, L., Arduino, M., Agüero, S. M., Gomez, P., et al. (1998). Outbreak of *Acinetobacter* spp. bloodstream infections in a nursery associated with contaminated aerosols and air conditioners. *Pediatric Infectious Disease Journal*, 17(8), 716-722.
- McKendrick, G. D., & Emond, R. T. (1976). Investigation of cross-infection in isolation wards of different design. *Journal of Hygiene*, 76(1), 23-31.
- McManus, A. T., Mason, A. D., Jr., McManus, W. F., & Pruitt, B. A., Jr. (1992). Control of *Pseudomonas aeruginosa* infections in burned patients. *Surgical Research Communications*, 12, 61-67.
- McManus, A. T., Mason, A. D., Jr., McManus, W. F., & Pruitt, B. A., Jr. (1994). A decade of reduced gram-negative infections and mortality associated with improved isolation of burned patients. *Archives of Surgery*, 129(12), 1306-1309.
- McManus, A. T., McManus, W. F., Mason, A. D., Jr., Aitcheson, A. R., & Pruitt, B. A., Jr. (1985). Microbial colonization in a new intensive care burn unit. A prospective cohort study. *Archives of Surgery*, 120(2), 217-223.
- McMurdo, M. E. T., Millar, A. M., & Daly, F. (2000). A randomized controlled trial of fall prevention strategies in old peoples' homes. *Gerontology*, 46(2), 83-87.
- Meengs, M. R., Giles, B. K., Chisholm, C. D., Cordell, W. H., & Nelson, D. R. (1994). Hand washing frequency in an emergency department. *Journal of Emergency Nursing*, 20(3), 183-188.
- Mehta, G. (1990). Aspergillus endocarditis after open heart surgery: An epidemiological investigation. *Journal of Hospital Infection*, 15(3), 245-253.
- Menzies, D., Fanning, A., Yuan, L., & FitzGerald, J. M. (2000). Hospital ventilation and risk for tuberculous infection in Canadian health care workers. *Annals of Internal Medicine*, 133(10), 779-789.
- Mermel, L. A., Josephson, S. L., Giorgio, C. H., Dempsey, J., & Parenteau, S. (1995). Association of Legionnaires' disease with construction: Contamination of potable water? *Infection Control and Hospital Epidemiology*, 16(2), 76-81.
- Merriman, E., Corwin, P., & Ikram, R. (2002). Toys are a potential source of cross-infection in general practitioners' waiting rooms. *British Journal of General Practice*, 52(475), 138-140.
- Mion, L. C., Fogel, J., Sandhu, S., Palmer, R. M., Minnick, A. F., Cranston, T., et al. (2001). Outcomes following physical restraint reduction programs in two acute care hospitals. *Joint Commission Journal on Quality Improvement*, 27(11), 605-618.
- Mlinek, E. J., & Pierce, J. (1997). Confidentiality and privacy breaches in a university hospital emergency department. *Academic Emergency Medicine*, 4(12), 1142-1146.
- Morawska, L., Jamriska, M., & Francis, P. (1998). Particulate matter in the hospital environment. *Indoor Air*, 8(4), 285-294.
- Morgan, V. R., Mathison, J. H., Rice, J. C., & Clemmer, D. I. (1985). Hospital falls: A persistent problem. *American Journal of Public Health*, 75(7), 775.
- Mortlock, T. (1994). Maintaining patient confidentiality in A&E. *Nursing Times*, 90(34), 42-43.

- Muto, C. A., Sistrom, M. G., & Farr, B. M. (2000). Hand hygiene rates unaffected by installation of dispensers of a rapidly acting hand antiseptic. *American Journal of Infection Control*, 28(3), 273-276.
- Nauseef, W. M., & Maki, D. G. (1981). A study of the value of simple protective isolation in patients with granulocytopenia. *New England Journal of Medicine*, 304(8), 448-453.
- Neely, A. N., & Maley, M. P. (2001). Dealing with contaminated computer keyboards and microbial survival. *American Journal of Infection Control*, 29(2), 131-132.
- Neely, A. N., & Sittig, D. F. (2002). Basic microbiologic and infection control information to reduce the potential transmission of pathogens to patients via computer hardware. *Journal of the American Medical Informatics Association*, 9(5), 500-508.
- Noskin, G. A., Bednarz, P., Suriano, T., Reiner, S., & Peterson, L. (2000). Persistent contamination of fabric-covered furniture by Vancomycin-resistant Enterococci: Implication for upholstery selection in hospitals. *American Journal of Infection Control*, 28(4), 311-313.
- Noskin, G. A., & Peterson, L. R. (2001). Engineering infection control through facility design. *Emerging Infectious Diseases*, 7(2), 354-357.
- Noskin, G. A., Stosor, V., Cooper, I., & Peterson, L. R. (1995). Recovery of vancomycin-resistant enterococci on fingertips and environmental surfaces. *Infection Control and Hospital Epidemiology*, 16(10), 577-581.
- Nurmi, I., & Luthje, P. (2002). Incidence and costs of falls and fall injuries among elderly in institutional care. *Scandinavian Journal of Primary Health Care*, 20(2), 118-122.
- Nystrom, B. (1983). Optimal design/personnel for control of intensive care unit infection. *Infection Control*, 4(5), 388-390.
- Nystrom, B. (1994). Impact of handwashing on mortality in intensive care: Examination of the evidence. *Infection Control and Hospital Epidemiology*, 15(7), 435-436.
- Obbard, J. P., & Fang, L. S. (2003). Airborne concentrations of bacteria in a hospital environment in Singapore. *Water Air and Soil Pollution*, 144(1), 333-341.
- O'Keefe, S. T. (2004). Down with bedrails? *Lancet*, 363(9406), 343-344.
- Opal, S. M., Asp, A. A., Cannady, P. B., Jr., Morse, P. L., Burton, L. J., & Hammer, P. G. (1986). Efficacy of infection control measures during a nosocomial outbreak of disseminated aspergillosis associated with hospital construction. *Journal of Infectious Diseases*, 153(3), 634-637.
- Oren, I., Haddad, N., Finkelstein, R., & Rowe, J. M. (2001). Invasive pulmonary aspergillosis in neutropenic patients during hospital construction: Before and after chemoprophylaxis and institution of HEPA filters. *American Journal of Hematology*, 66(4), 257-262.
- Palmer, R. (1999). Bacterial contamination of curtains in clinical areas. *Nursing Standard*, 14(2), 33-35.
- Panagopoulou, P., Filioti, J., Petrikos, G., Giakouppi, P., Anatoliotaki, M., Farmaki, E., et al. (2002). Environmental surveillance of filamentous fungi in three tertiary care hospitals in Greece. *Journal of Hospital Infection*, 52(3), 185-191.
- Parker, K., & Miles, S. H. (1997). Deaths caused by bedrails. *Journal of the American Geriatrics Society*, 45(7), 797-802.
- Passweg, J. R., Rowlings, P. A., Atkinson, K. A., Barrett, A. J., Gale, R. P., Gratwohl, A., et al. (1998). Influence of protective isolation on outcome of allogeneic bone marrow transplantation for leukemia. *Bone Marrow Transplantation*, 21(12), 1231-1238.
- Pegues, D. A., & Woernle, C. H. (1993). An outbreak of acute nonbacterial gastroenteritis in a nursing home. *Infection Control and Hospital Epidemiology*, 14(2), 87-94.
- Pettinger, A., & Nettleman, M. D. (1991). Epidemiology of isolation precautions. *Infection Control and Hospital Epidemiology*, 12(5), 303-307.
- Pittet, D., Hugonnet, S., Harbarth, S., Mourouga, P., Sauvan, V., Touveneau, S., et al. (2000).

- Effectiveness of a hospital-wide programme to improve compliance with hand hygiene. *Lancet*, 356(9238), 1307-1312.
- Pittet, D., Mourouga, P., & Perneger, T. V. (1999). Compliance with handwashing in a teaching hospital. *Annals of Internal Medicine*, 130(2), 126-130
- Preston, G. A., Larson, E. L., & Stamm, W. E. (1981). The effect of private isolation rooms on patient care practices, colonization and infection in an intensive care unit. *American Journal of Medicine*, 70(3), 641-645.
- Pryor, F., & Messmer, P. R. (1998). The effect of traffic patterns in the OR on surgical site infections. *AORN Journal*, 68(4), 649-660.
- Raad, I., Hanna, H., Osting, C., Hachem, R., Umphrey, J., Tarrand, J., et al. (2002). Masking of neutropenic patients on transport from hospital rooms is associated with a decrease in nosocomial aspergillosis during construction. *Infection Control and Hospital Epidemiology*, 23(1), 41-43.
- Redman, J. F., & McNatt, S. J. (2000). Portable cushioned operating table siderails: An adjunct to pediatric surgery. *Southern Medical Journal*, 93(11), 1081-1082.
- Reiling, J., Breckbill, C., Murphy, M., McCullough, S., & Chernos, S. (2003). Facility designing around patient safety and its effect on nursing. *Nursing Economics*, 21(3), 143-147.
- Resnick, B. (1999). Falls in a community of older adults: Putting research into practice. *Clinical Nursing Research*, 8(3), 251-266.
- Rice, N., Streifel, A., & Vesley, D. (2001). An evaluation of hospital special-ventilation-room pressures. *Infection Control and Hospital Epidemiology*, 22(1), 19-23.
- Roberts, S. A., Findlay, R., & Lang, S. D. (2001). Investigation of an outbreak of multi-drug resistant *Acinetobacter baumannii* in an intensive care burns unit. *Journal of Hospital Infection*, 48(3), 228-232.
- Rolandi, L., Lodola, L., Guglielminetti, M., Caretta, G., & Azzaretti, G. (1998). Evaluation of airborne particulate and fungi in critical hospital care units. *Toxicology Letters*, 95(Suppl. 1), 226.
- Roseman, C., & Booker, J. M. (1995). Workload and environmental factors in hospital medication errors. *Nursing Research*, 44(4), 226-230.
- Rountree, P. M., Beard, M. A., Loewenthal, J., May, J., & Renwick, S. B. (1967). Staphylococcal sepsis in a new surgical ward. *British Medical Journal*, 1(533), 132-137.
- Saiman, L. (2002). Risk factors for hospital-acquired infections in the neonatal intensive care unit. *Seminars in Perinatology*, 26(5), 315-321.
- Samore, M. H., Venkataraman, L., DeGirolami, P. C., Arbeit, R. D., & Karchmer, A. W. (1996). Clinical and molecular epidemiology of sporadic and clustered cases of nosocomial *Clostridium difficile* diarrhea. *American Journal of Medicine*, 100(1), 32-40.
- San Jose-Alonso, J. F., Velasco-Gomez, E., Rey-Martinez, F. J., Alvarez-Guerra, M., & Pelaez, C. G. (1999). Study on environmental quality of a surgical block. *Energy and Buildings*, 29(2), 179-187.
- Sanderson, P. J., & Weissler, S. (1992). Recovery of coliforms from the hands of nurses and patients: Activities leading to contamination. *Journal of Hospital Infection*, 21(2), 85-93.
- Sarubbi, F. A., Jr., Kopf, H. B., Wilson, M. B., McGinnis, M. R., & Rutala, W. A. (1982). Increased recovery of *Aspergillus flavus* from respiratory specimens during hospital construction. *American Review of Respiratory Disease*, 125(1), 33-38.
- Schott, J., & Force, R. W. (1999). Do bedrails decrease falls in hospitalized patients? *Journal of Family Practice*, 48(8), 579.
- Sehulster, L., & Chinn, R. Y. (2003). Guidelines for environmental infection control in health-care facilities. Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee (HICPAC). *MMWR. Recommendations and Reports*, 52(RR-10), 1-42.



- Sheperd, L., & Foureur, M. (2001). Introduction of night-lights to reduce falls on an assessment, treatment and rehabilitation unit. *JARNA*, 4(4), 11-13.
- Sherertz, R., J. Belani, A., Kramer, B. S., Elfenbein, G. J., Weiner, R. S., Sullivan, M. L., et al. (1987). Impact of air filtration on nosocomial Aspergillus infections: Unique risk of bone marrow transplant recipients. *American Journal of Medicine*, 83(4), 709-718.
- Sherertz, R. J., & Sullivan, M. L. (1985). An outbreak of infections with Acinetobacter calcoaceticus in burn patients: Contamination of patients' mattresses. *Journal of Infectious Diseases*, 151(2), 252-258.
- Shirani, K. Z., McManus, A. T., Vaughan, G. M., McManus, W. F., Pruitt, B. A., Jr., & Mason, A. D., Jr. (1986). Effects of environment on infection in burn patients. *Archives of Surgery*, 121(1), 31-36.
- Skoutelis, A. T., Westenfelder, G. O., Beckerdite, M., & Phair, J. P. (1994). Hospital carpeting and epidemiology of Clostridium difficile. *American Journal of Infection Control*, 22(4), 212-217.
- Smylie, H. G., Davidson, A. I., Macdonald, A., & Smith, G. (1971). Ward design in relation to postoperative wound infection. *British Medical Journal*, 1(740), 67-72.
- Soe, J. A. T., Hall, J. M., Collyns, T. A., Witherden, I. R., Parnell, P., Woodrow, G., et al. (2002). An outbreak of vancomycin-resistant enterococci associated with major ward refurbishment. *Journal of Hospital Infection*, 50(1), 79-80.
- Srinivasan, A., Beck, C., Buckley, T., Geyh, A., Bova, G., Merz, W., et al. (2002). The ability of hospital ventilation systems to filter Aspergillus and other fungi following a building implosion. *Infection Control and Hospital Epidemiology*, 23(9), 520-524.
- Startzell, J. K., Owens, D. A., Mulfinger, L. M., & Cavanagh, P. R. (2000). Stair negotiation in older people: A review. *Journal of the American Geriatrics Society*, 48(5), 567-580.
- Streifel, A. J., Vesley, D., Rhame, F. S., & Murray, B. (1989). Control of airborne environmental spores in a university hospital. *Environment International*, 15(1-6), 221-227.
- Stuyver, L., Claeys, H., Wyseur, A., Van Arnhem, W., De Beenhouwer, H., Uytendaele, S., et al. (1996). Hepatitis C virus in a hemodialysis unit: Molecular evidence for nosocomial transmission. *Kidney International*, 49(3), 889-895.
- Su, L., Li, H., & Bai, T. (1996). The design of air purification system and its effect on surgical ICU. *Zhonghua Wai Ke Za Zhi [Chinese Journal of Surgery]*, 34(5), 291-293.
- Tabbara, K. F., & Al Jabarti, A. (1998). Hospital construction-associated outbreak of ocular aspergillosis after cataract surgery. *Ophthalmology*, 105(3), 522-526.
- Talon, D., Vichard, P., Muller, A., Bertin, M., Jeunet, L., & Bertrand, X. (2003). Modelling the usefulness of a dedicated cohort facility to prevent the dissemination of MRSA. *Journal of Hospital Infection*, 54(1), 57-62.
- Thompson, J. T., Meredith, J. W., & Molnar, J. A. (2002). The effect of burn nursing units on burn wound infections. *Journal of Burn Care & Rehabilitation*, 23(4), 281-286; discussion 280.
- Ubel, P. A., Zell, M. M., & Miller, D. J. (1995). Elevator talk: Observational study of inappropriate comments in a public space. *American Journal of Medicine*, 99(2), 190-194.
- Uden, G. (1985). Inpatient accidents in hospitals. *Journal of the American Geriatrics Society*, 33(12), 833-841.
- Uduman, S. A., Farrukh, A. S., Nath, K. N., Zuhair, M. Y., Ifrah, A., Khawla, A. D., et al. (2002). An outbreak of Serratia marcescens infection in a special-care baby unit of a community hospital in United Arab Emirates: The importance of the air conditioner duct as a nosocomial reservoir. *Journal of Hospital Infection*, 52(3), 175-180.
- Lee, M. L., Chen, C. J., Su, I., Chen, K. T., Yeh, C. C., King, C. C., et al. (2003). Use of quarantine to prevent transmission of severe acute respiratory syndrome—Taiwan, 2003.

- MMWR*, 52(29), 680-683.
- Utrup, L. J., Werner, K., & Frey, A. H. (2003). Minimizing pathogenic bacteria, including spores, in indoor air. *Journal of Environmental Health*, 66(5), 19-26, 29.
- van Leeuwen, M., Bennett, L., West, S., Wiles, V., & Grasso, J. (2001). Patient falls from bed and the role of bedrails in the acute care setting. *Australian Journal of Advanced Nursing*, 19(2), 8-13.
- Vassallo, M., Azeem, T., Pirwani, M. F., Sharma, J. C., & Allen, S. C. (2000). An epidemiological study of falls on integrated general medical wards. *International Journal of Clinical Practice*, 54(10), 654-657.
- Vernon, M. O., Trick, W. E., Welbel, S. F., Peterson, B. J., & Weinstein, R. A. (2003). Adherence with hand hygiene: Does number of sinks matter? *Infection Control and Hospital Epidemiology*, 24(3), 224-225.
- Volpp, K. G. M., & Grande, D. (2003). Residents' suggestions for reducing errors in teaching hospitals. *New England Journal of Medicine*, 348(9), 851-855.
- Voss, A., & Widmer, A. F. (1997). No time for handwashing? Handwashing versus alcoholic rub: Can we afford 100% compliance? *Infection Control and Hospital Epidemiology*, 18(3), 205-208.
- Walsh, T. R., Guttendorf, J., Dummer, S., Hardesty, R. L., Armitage, J. M., Kormos, R. L., et al. (1989). The value of protective isolation procedures in cardiac allograft recipients. *Annals of Thoracic Surgery*, 47(4), 539-544; discussion 544-535.
- Wan, G. H., Lu, S. C., & Tsai, Y. H. (2004). Polymerase chain reaction used for the detection of airborne Mycobacterium tuberculosis in health care settings. *American Journal of Infection Control*, 32(1), 17-22.
- Ward, A., Candela, L., & Mahoney, J. (2004). Developing a unit-specific falls reduction program. *Journal for Healthcare Quality*, 26(2), 36-40; quiz 40-31.
- Watanakunakorn, C., Wang, C., & Hazy, J. (1998). An observational study of hand washing and infection control practices by healthcare workers. *Infection Control and Hospital Epidemiology*, 19(11), 858-860.
- Weems, J. J., Jr., Davis, B. J., Tablan, O. C., Kaufman, L., & Martone, W. J. (1987). Construction activity: An independent risk factor for invasive aspergillosis and zygomycosis in patients with hematologic malignancy. *Infection Control*, 8(2), 71-75.
- Wendt, C., Dietze, B., Dietz, E., & Ruden, H. (1997). Survival of *Acinetobacter baumannii* on dry surfaces. *Journal of Clinical Microbiology*, 35(6), 1394-1397.
- Wenzel, R. P., Thompson, R. L., Landry, S. M., Russell, B. S., Miller, P. J., Ponce de Leon, S., et al. (1983). Hospital-acquired infections in intensive care unit patients: An overview with emphasis on epidemics. *Infection Control*, 4(5), 371-375.
- Wigglesworth, N. (2003). The use of protective isolation. *Nursing Times*, 99(7), 26-27.
- Williams, H. N., Singh, R., & Romberg, E. (2003). Surface contamination in the dental operator: A comparison over two decades. *Journal of the American Dental Association*, 134(3), 325-330; quiz 339.
- Wolf, E. J. (2003). Promoting patient safety through facility design. *Healthcare Executive*, 18(4), 16-20.
- Wong, S., Glennie, K., Muise, M., Lambie, E., & Meagher, D. (1981). An exploration of environmental variables and patient falls. *Dimensions in Health Service*, 58(6), 9-11.
- Wu, P. C., Su, H. J., & Ho, H. M. (2000). A comparison of sampling media for environmental viable fungi collected in a hospital environment. *Environmental Research*, 82(3), 253-257.
- Zimmerman, S., Gruber-Baldini, A. L., Hebel, J. R., Sloane, P. D., & Magaziner, J. (2002). Nursing home facility risk factors for infection and hospitalization: Importance of registered nurse turnover, administration, and social factors. *Journal of the American*

*Geriatrics Society*, 50(12), 1987-1995.

***Reduce Stress and Improve Outcomes***

- Aaron, J. N., Carlisle, C. C., Carskadon, M. A., Meyer, T. J., Hill, N. S., & Millman, R. P. (1996). Environmental noise as a cause of sleep disruption in an intermediate respiratory care unit. *Sleep*, 19(9), 707-710.
- Ackerman, B., Sherwonit, E., & Fisk, W. (1989). Reduced incidental light exposure: Effect on the development of retinopathy of prematurity in low birth weight infants. *Pediatrics*, 83(6), 958-962.
- Aitken, R. J. (1982). Quantitative noise analysis in a modern hospital. *Archives of Environmental Health*, 37(6), 361-364.
- Alessi, C. A., Schnelle, J. F., MacRae, P. G., Ouslander, J. G., al-Samarrai, N., Simmons, S. F., et al. (1995). Does physical activity improve sleep in impaired nursing home residents? *Journal of the American Geriatrics Society*, 43(10), 1098-1102.
- Allaouchiche, B., Duflo, F., Debon, R., Bergeret, A., & Chassard, D. (2002). Noise in the postanaesthesia care unit. *British Journal of Anaesthesia*, 88(3), 369-373.
- Anagnostakis, D., Petmezakis, J., Messaritakis, J., & Matsaniotis, N. (1980). Noise pollution in neonatal units: A potential health hazard. *Acta Paediatrica Scandinavica*, 69(6), 771-773.
- Astedt-Kurki, P., Paunonen, M., & Lehti, K. (1997). Family members' experiences of their role in a hospital: A pilot study. *Journal of Advanced Nursing*, 25(5), 908-914.
- Avery, D. H. (1998). A turning point for seasonal affective disorder and light therapy research? *Archives of General Psychiatry*, 55(10), 863.
- Baker, C. F. (1984). Sensory overload and noise in the ICU: Sources of environmental stress. *Critical Care Quarterly*, 6(4), 66-80.
- Baker, C. F. (1992). Discomfort to environmental noise: Heart rate responses of SICU patients. *Critical Care Nursing Quarterly*, 15(2), 75-90.
- Baker, C. F. (1993). Annoyance to ICU noise: A model of patient discomfort. *Critical Care Nursing Quarterly*, 16(2), 83-90.
- Baker, C. F., Garvin, B. J., Kennedy, C. W., & Polivka, B. J. (1993). The effect of environmental sound and communication on CCU patients' heart rate and blood pressure. *Research in Nursing & Health*, 16(6), 415-421.
- Baldwin, S. (1985). Effects of furniture rearrangement on the atmosphere of wards in a maximum-security hospital. *Hospital & Community Psychiatry*, 36(5), 525-528.
- Balogh, D., Kittinger, E., Benzer, A., & Hackl, J. M. (1993). Noise in the ICU. *Intensive Care Medicine*, 19(6), 343-346.
- Barnhart, S. K., Perkins, N. H., & Fitzsimonds, J. (1998). Behaviour and outdoor setting preferences at a psychiatric hospital. *Landscape and Urban Planning*, 42(2), 147-156.
- Barry, J. T., & Miller, D. B. (1980). The nursing home visitor: Who, when, where and for how long? *Long Term Care and Health Services Administration Quarterly*, 4(4), 261-274.
- Bay, E. J., Kupferschmidt, B., Opperwall, B. J., & Speer, J. (1988). Effect of the family visit on the patient's mental status. *Focus on Critical Care*, 15(1), 11-16.
- Bayo, M. V., Garcia, A. M., & Garcia, A. (1995). Noise levels in an urban hospital and workers' subjective responses. *Archives of Environmental Health*, 50(3), 247-251.
- Beauchemin, K. M., & Hays, P. (1996). Sunny hospital rooms expedite recovery from severe and refractory depressions. *Journal of Affective Disorders*, 40(1), 49-51.
- Beauchemin, K. M., & Hays, P. (1998). Dying in the dark: Sunshine, gender and outcomes in myocardial infarction. *Journal of the Royal Society of Medicine*, 91(7), 352-354.
- Bell, P. L. (1997). Adolescent mothers' perceptions of the neonatal intensive care unit

- environment. *Journal of Perinatal & Neonatal Nursing*, 11(1), 77-84.
- Belliemi, C. V., Buonocore, G., Pinto, I., Stacchini, N., Cordelli, D. M., & Bagnoli, F. (2003). Use of sound-absorbing panel to reduce noisy incubator reverberating effects. *Biology of the Neonate*, 84(4), 293-296.
- Benedetti, F., Colombo, C., Barbini, B., Campori, E., & Smeraldi, E. (2001). Morning sunlight reduces length of hospitalization in bipolar depression. *Journal of Affective Disorders*, 62(3), 221-223.
- Benians, R. C. (1988). The influence of parental visiting on survival and recovery of extensively burned children. *Burns, Including Thermal Injury*, 14(1), 31-34.
- Benini, F., Magnavita, V., Lago, P., Arslan, E., & Pisan, P. (1996). Evaluation of noise in the neonatal intensive care unit. *American Journal of Perinatology*, 13(1), 37-41.
- Bentley, S., Murphy, F., & Dudley, H. (1977). Perceived noise in surgical wards and an intensive care area: An objective analysis. *British Medical Journal*, 2(6101), 1503-1506.
- Berens, R. J., & Weigle, C. G. (1996). Cost analysis of ceiling tile replacement for noise abatement. *Journal of Perinatology*, 16(Pt 1), 199-201.
- Berg, S. (2001). Impact of reduced reverberation time on sound-induced arousals during sleep. *Sleep*, 24(3), 289-292.
- Berglund, B., Lindvall, T., & Schwela, D. H. (1999). *Guidelines for community noise*. World Health Organization: Protection of the Human Environment.
- Berrutto, V., Fontoynt, M., & Gue, J. M. F. (1997). Effect of temperature and light source type (natural/artificial) on visual comfort appraisal: Experimental design and setting. *Right Light 4 Proceedings*, 1, 139-141.
- Berwick, D. M. & Kotagal, M. (2004). Restricted visiting hours in ICUs: Time to change. *JAMA*, 292(6), 736-737.
- Biley, F. C. (1994). Effects of noise in hospitals. *British Journal of Nursing*, 3(3), 110-113.
- Blackburn, S., & Patteson, D. (1991). Effects of cycled light on activity state and cardiorespiratory function in preterm infants. *Journal of Perinatal & Neonatal Nursing*, 4(4), 47-54.
- Blomkvist, V., Eriksen, C. A., Theorell, T., Ulrich, R. S., & Rasmanis, G. (in press). Acoustics and psychosocial environment in coronary intensive care. *Occupational and Environmental Medicine*.
- Bonn, D. (1998). Noise pollution in intensive-care units reduced by simple measures. *Lancet*, 352(9128), 628.
- Boubekri, M., Hull, R. B., & Boyer, L. L. (1991). Impact of window size and sunlight penetration on office workers' mood and satisfaction: A novel way of assessing sunlight. *Environment & Behavior*, 23(4), 474-493.
- Bovenzi, M., & Collareta, A. (1984). Noise levels in a hospital. *Industrial Health*, 22(2), 75-82.
- Bremmer, P., Byers, J. F., & Kiehl, E. (2003). Noise and the premature infant: Physiological effects and practice implications. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 32(4), 447-454.
- Brown, B., Wright, H., & Brown, C. (1997). A post-occupancy evaluation of wayfinding in a pediatric hospital: Research findings and implications for instruction. *Journal of Architectural and Planning Research*, 14(1), 35-51.
- Brown, P., & Taquino, L. T. (2001). Designing and delivering neonatal care in single rooms. *Journal of Perinatal & Neonatal Nursing*, 15(1), 68-83.
- Buemi, M., Allegra, A., Grasso, F., & Mondio, G. (1995). Noise pollution in an intensive care unit for nephrology and dialysis. *Nephrology, Dialysis, Transplantation*, 10(12), 2235-2239.
- Butler, D., Acquino, A. L., Hissong, A. A., & Scott, P. A. (1993). Wayfinding by newcomers in a complex building. *Human Factors*, 25(1), 159-173.

- Callahan, E. J., Brasted, W. S., Myerberg, D. Z., & Hamilton, S. (1991). Prolonged travel time to neonatal intensive care unit does not affect content of parental visiting: A controlled prospective study. *Journal of Rural Health* 7(1), 73-83.
- Carpman, J. R. (1993). *Design that cares: Planning health facilities for patients and visitors* (2nd ed.). Chicago: American Hospital Publishing.
- Carpman, J. R., Grant, M., & Simmons, D. (1983-84). Wayfinding in the hospital environment: The impact of various floor numbering alternatives. *Journal of Environmental Systems*, 13(4), 353-364.
- Carpman, J. R., Grant, M., & Simmons, D. (1984). *No more mazes: Research about design for wayfinding in hospitals*. Ann Arbor, MI: The University of Michigan Hospitals.
- Carpman, J. R., Grant, M. A., & Simmons, D. A. (1985). Hospital design and wayfinding: A video simulation study. *Environment & Behavior*, 17(3), 296-314.
- Carr, J. A., & Powers, M. J. (1986). Stressors associated with coronary bypass surgery. *Nursing Research*, 35(4), 243-246.
- Causey, D. L., McKay, M., Rosenthal, C., & Darnell, C. (1998). Assessment of hospital-related stress in children and adolescents admitted to a psychiatric inpatient unit. *Journal of Child and Adolescent Psychiatric Nursing*, 11(4), 135-145.
- Cederberg, R. A., Frederiksen, N. L., Benson, B. W., & Shulman, J. D. (1998). Effect of different background lighting conditions on diagnostic performance of digital and film images. *Dento Maxillo Facial Radiology*, 27(5), 293-297.
- Chang, Y. J., Lin, C. H., & Lin, L. H. (2001). Noise and related events in a neonatal intensive care unit. *Acta Paediatrica Taiwanica = Taiwan Er Ke Yi Xue Hui Za Zhi*, 42(4), 212-217.
- Cheek, F. E., Maxwell, R., & Weisman, R. (1971). Carpeting the ward: An exploratory study in environmental psychiatry. *Mental Hygiene*, 55(1), 109-118.
- Chen, H. F., & Chang, Y. J. (2001). Noise distribution of an incubator with nebulizer at a neonatal intensive care unit in southern Taiwan. *Journal of Nursing Research*, 9(3), 25-32.
- Christenfeld, R., Wagner, J., Pastva, G., & Acrish, W. P. (1989). How physical settings affect chronic mental patients. *Psychiatric Quarterly*, 60(3), 253-264.
- Cmiel, C. A., Karr, D. M., Gasser, D. M., Oliphant, L. M., & Neveau, A. J. (2004). Noise control: A nursing team's approach to sleep promotion. *American Journal of Nursing*, 104(2), 40-48.
- Cohen-Mansfield, J., & Werner, P. (1999). Outdoor wandering parks for persons with dementia: A survey of characteristics and use. *Alzheimer's Disease and Associated Disorders*, 13(2), 109-117.
- Colledge, J. (1980). Visiting the mentally handicapped in residential care. *British Journal of Psychiatry*, 137, 313-318.
- Cooper Marcus, C. (2003). Healing havens: Two hospital gardens in Portland, Oregon, win awards for therapeutic values. *Landscape Architecture*, 93(8): 84, 91, 107-109.
- Cooper Marcus, C., & Barnes, M. (1995). *Gardens in healthcare facilities: Uses, therapeutic benefits, and design recommendations*. Martinez, CA: Center for Health Design.
- Corser, N. C. (1996). Sleep of 1- and 2-year-old children in intensive care. *Issues in Comprehensive Pediatric Nursing*, 19(1), 17-31.
- Couper, R. T., Hendy, K., Lloyd, N., Gray, N., Williams, S., & Bates, D. J. (1994). Traffic and noise in children's wards. *Medical Journal of Australia*, 160(6), 338-341.
- Cropp, A. J., Woods, L. A., Raney, D., & Bredle, D. L. (1994). Name that tone. The proliferation of alarms in the intensive care unit. *Chest*, 105(4), 1217-1220.
- Cureton-Lane, R. A., & Fontaine, D. K. (1997). Sleep in the pediatric ICU: An empirical investigation. *American Journal of Critical Care*, 6(1), 56-63.

- Cuttini, M., Rebagliato, M., Bortoli, P., Hansen, G., de Leeuw, R., Lenoir, S., et al. (1999). Parental visiting, communication, and participation in ethical decisions: A comparison of neonatal unit policies in Europe. *Archives of Disease in Childhood. Fetal and Neonatal Edition*, 81(2), F84-91.
- Daniels, D. Y. (1996). Visiting hours. An open ICU. *RN*, 59(10), 30-32.
- Davies, J. M., Ewen, A., Cuppage, A., Gilbert, D., & Winkelaar, R. (1989). Noise levels in operating rooms—A comparison of Canada and England. *Anaesthesia and Intensive Care*, 17(1), 98.
- Deep, P., & Petropoulos, D. (2003). Effect of illumination on the accuracy of identifying interproximal carious lesions on bitewing radiographs. *Journal (Canadian Dental Association)*, 69(7), 444-446.
- Diette, G. B., Lechtzin, N., Haponik, E., Devrotes, A., & Rubin, H. R. (2003). Distraction therapy with nature sights and sounds reduces pain during flexible bronchoscopy: A complementary approach to routine analgesia. *Chest*, 123(3), 941-948.
- Diffey, B. L., & Storey, A. (1988). Light and length of stay in hospital. *Journal of the Royal Society of Medicine*, 81(11), 643.
- Dijkers, M., Yavuzer, G., Ergin, S., Weitzenkamp, D., & Whiteneck, G. G. (2002). A tale of two countries: Environmental impacts on social participation after spinal cord injury. *Spinal Cord*, 40(7), 351-362.
- Dodenhoff, R. M. (1995). Noise in the orthopaedic operating theatre. *Annals of the Royal College of Surgeons of England*, 77(Suppl.), 8-9.
- Donchin, Y., & Seagull, F. J. (2002). The hostile environment of the intensive care unit. *Current Opinion in Critical Care*, 8(4), 316-320.
- Drinkard, J. L. (1984). *Wayfinding in the hospital environment: A design analysis*. Unpublished master's thesis, Georgia Institute of Technology, Atlanta.
- Duxbury, J. (1994). Avoiding disturbed sleep in hospitals. *Nursing Standard*, 9(10), 31-34.
- Dvoskin, J. A., Radomski, S. J., Bennett, C., Olin, J. A., Hawkins, R. L., Dotson, L. A., et al. (2002). Architectural design of a secure forensic state psychiatric hospital. *Behavioral Sciences & the Law*, 20(5), 481-493.
- Eastman, C. I., Young, M. A., Fogg, L. F., Liu, L., & Meaden, P. M. (1998). Bright light treatment of winter depression. *Archives of General Psychiatry*, 55(10), 883.
- Elander, G., & Hellstrom, G. (1995). Reduction of noise levels in intensive care units for infants: Evaluation of an intervention program. *Heart & Lung*, 24(5), 376-379.
- Epp, T., MacPhee, R., & Allman, A. (2001). Let there be light: The aging eye needs more illumination than is available in many facilities. *Canadian Nursing Home*, 12(2), 4.
- Falk, S. A., & Woods, N. F. (1973). Hospital noise-levels and potential health hazards. *New England Journal of Medicine*, 289(15), 774-781.
- Federman, E. J., Drebing, C. E., Boisvert, C., Penk, W., Binus, G. & Rosenheck, R. (2000). Relationship between climate and psychiatric inpatient length of stay in Veterans Health Administration hospitals. *American Journal of Psychiatry*, 157(10), 1669.
- Fife, D., & Rappaport, E. (1976). Noise and hospital stay. *American Journal of Public Health*, 66(7), 680-681.
- Finlay, G. (1991). Sleep and intensive care. *Intensive Care Nursing*, 7(1), 61-68.
- Firestone, I. J., Lichtman, C. M., & Evans, J. R. (1980). Privacy and solidarity: Effects of nursing home accommodation on environmental perception and sociability preferences. *International Journal of Aging & Human Development*, 11(3), 229-241.
- Flynn, E. A., Barker, K. N., Gibson, J. T., Pearson, R. E., Smith, L. A., & Berger, B. A. (1996). Relationships between ambient sounds and the accuracy of pharmacists' prescription-filling performance. *Human Factors*, 38(4), 614-622.
- Flynn, M. B. (1996). Wound healing and critical illness. *Critical Care Nursing Clinics of North*

- America*, 8(2), 115-123.
- Foltz-Gray, D. (1995). Intimate strangers. When roommates clash, caregivers can ease the tension—or make it worse. *Contemporary Longterm Care*, 18(6), 34-37.
- Foss, K. R., & Tenholder, M. F. (1993). Expectations and needs of persons with family members in an intensive care unit as opposed to a general ward. *Southern Medical Journal*, 86(4), 380-384.
- Fox, R. A., & Henson, P. W. (1996). Potential ocular hazard from a surgical light source. *Australasian Physical and Engineering Sciences in Medicine*, 19(1), 12-16.
- Franck, L. S., & Spencer, C. (2003). Parent visiting and participation in infant caregiving activities in a neonatal unit. *Birth*, 30(1), 31-35.
- Freedman, N. S., Gazendam, J., Levan, L., Pack, A. I., & Schwab, R. J. (2001). Abnormal sleep/wake cycles and the effect of environmental noise on sleep disruption in the intensive care unit. *American Journal of Respiratory and Critical Care Medicine*, 163(2), 451-457.
- Freedman, N. S., Kotzer, N., & Schwab, R. J. (1999). Patient perception of sleep quality and etiology of sleep disruption in the intensive care unit. *American Journal of Respiratory and Critical Care Medicine*, 159(Pt 1), 1155-1162.
- Friberg, T. R., & Borrero, G. (2000). Diminished perception of ambient light: A symptom of clinical depression? *Journal of Affective Disorders*, 61(1-2), 113-118.
- Frumkin, H. (2001). Beyond toxicity: Human health and the natural environment. *American Journal of Preventive Medicine*, 20(3), 234-240.
- Gabor, J. Y., Cooper, A. B., Crombach, S. A., Lee, B., Kadikar, N., Bettger, H. E., et al. (2003). Contribution of the intensive care unit environment to sleep disruption in mechanically ventilated patients and healthy subjects. *American Journal of Respiratory and Critical Care Medicine*, 167(5), 708-715.
- Gast, P. L., & Baker, C. F. (1989). The CCU patient: Anxiety and annoyance to noise. *Critical Care Nursing Quarterly*, 12(3), 39-54.
- Gershon, J., Zimand, E., Lemos, R., Rothbaum, B. O., & Hodges, L. (2003). Use of virtual reality as a distractor for painful procedures in a patient with pediatric cancer: A case study. *Cyberpsychology & Behavior*, 6(6), 657-661.
- Giacoaia, G. P., Rutledge, D., & West, K. (1985). Factors affecting visitation of sick newborns. *Clinical Pediatrics*, 24(5), 259-262.
- Giunta, F., & Rath, J. (1969). Effect of environmental illumination in prevention of hyperbilirubinemia of prematurity. *Pediatrics*, 44(2), 162-167.
- Glass, P., Avery, G., Subramaniam, K. N. S., Keys, M. P., Sostek, A. M., & Friendly, D. S. (1985). Effect of bright light in the hospital nursery on the incidence of retinopathy of maturity. *New England Journal of Medicine*, 313(7), 401-404.
- Glod, C. A., Teicher, M. H., Butler, M., Savino, M., Harper, D., Magnus, E., et al. (1994). Modifying quiet room design enhances calming of children and adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 33(4), 558-566.
- Grant, M. (1988). "Wayfinding" in Michigan hospitals. *Michigan Hospitals*, 24(5), 45-47.
- Griffin, J. P., Myers, S., Kopelke, C., & Walker, D. (1988). The effects of progressive muscular relaxation on subjectively reported disturbance due to hospital noise. *Behavioral Medicine*, 14(1), 37-42.
- Grover, P. (1971). *Wayfinding in hospital environments: UCLA hospital disorientation pilot case study*. Los Angeles: Graduate School of Architecture and Urban Planning, University of California.
- Grumet, G. W. (1993). Pandemonium in the modern hospital. *New England Journal of Medicine*, 328(6), 433-437.
- Grumet, G. W. (1994). Noise hampers healing and curbs productivity. *Health Facilities*

- Management*, 7(1), 22-25.
- Haas, E. C., & Edworthy, J. (1996). Designing urgency into auditory warnings using pitch, speed and loudness. *Computing & Control Engineering Journal*, 7(4), 193-198.
- Haddock, J. (1994). Reducing the effects of noise in hospital. *Nursing Standard* 8(43), 25-28.
- Hagerman, I., Theorell, T., Ulrich, R. S., Blomkvist, V., Eriksen, C. A., & Rasmanis, G. (in press). Influence of coronary intensive care acoustics on the physiological states and quality of care of patients. *International Journal of Cardiology*.
- Haq, S. (2003). Investigating the syntax line: Configurational properties and cognitive correlates. *Environment & Planning B: Planning & Design*, 30, 841-863.
- Haq, S. (2001). Complex architectural settings: An investigation of spatial and cognitive variables through wayfinding behavior. *Dissertation Abstracts International Section A: Humanities & Social Sciences*, 62(2), 362.
- Haq, S. & Zimring, C. (2003). Just down the road a piece: The development of topological knowledge of building layouts. *Environment & Behavior*, 35(1), 132-160.
- Harris, D. (2000). *Environmental quality and healing environments: A study of flooring materials in a healthcare telemetry unit*. Doctoral dissertation, Texas A&M University, College Station.
- Harris, P. B., McBride, G., Ross, C., & Curtis, L. (2002). A place to heal: Environmental sources of satisfaction among hospital patients. *Journal of Applied Social Psychology*, 32(6), 1276-1299.
- Harris, R. W., & Reitz, M. L. (1985). Effects of room reverberation and noise on speech discrimination by the elderly. *Audiology*, 24(5), 319-324.
- Haslam, P. (1970). Caring for the total patient. Noise in hospitals: Its effect on the patient. *The Nursing Clinics of North America*, 5(4), 715-724.
- Heath, Y., & Gifford, R. (2001). Post-occupancy evaluation of therapeutic gardens in a multi-level care facility for the aged. *Activities, Adaptation & Aging*, 25(2), 21-43.
- Hebert, M., Dumont, M., & Paquet, J. (1998). Seasonal and diurnal patterns of human illumination under natural conditions. *Chronobiology International*, 15(1), 59-70.
- Hefferman, M. L., Morstatt, M., Saltzman, K., & Strunc, L. (1995). *A room with a view art survey: The Bird Garden at Duke University Hospital*. Durham, NC: Cultural Services Program and Management Fellows Program, Duke University Medical Center.
- Hilton, A. (1987). The hospital racket: How noisy is your unit? *American Journal of Nursing*, 87(1), 59-61.
- Hilton, B. A. (1976). Quantity and quality of patients' sleep and sleep-disturbing factors in a respiratory intensive care unit. *Journal of Advanced Nursing*, 1(6), 453-468.
- Hilton, B. A. (1985). Noise in acute patient care areas. *Research in Nursing & Health*, 8(3), 283-291.
- Hodge, B., & Thompson, J. F. (1990). Noise pollution in the operating theatre. *Lancet*, 335(8694), 891-894.
- Hoffman, H. G., Doctor, J. N., Patterson, D. R., Carrougher, G. J., & Furness, T. A., 3rd. (2000). Virtual reality as an adjunctive pain control during burn wound care in adolescent patients. *Pain*, 85(1-2), 305-309.
- Holahan, C. J. (1972). Seating patterns and patient behavior in an experimental dayroom. *Journal of Abnormal Psychology*, 80(2), 115-124.
- Holahan, C. J. (1976). Environmental change in a psychiatric setting: A social systems analysis. *Human Relations*, 29(2), 153-166.
- Holmberg, S. K., & Coon, S. (1999). Ambient sound levels in a state psychiatric hospital. *Archives of Psychiatric Nursing*, 13(3), 117-126.
- Hook, W. F., Sobal, J., & Oak, J. C. (1982). Frequency of visitation in nursing homes: Patterns of contact across the boundaries of total institutions. *Gerontologist*, 22(4), 424-428.



- Hutton, A. (2002). The private adolescent: Privacy needs of adolescents in hospitals. *Journal of Pediatric Nursing, 17*(1), 67-72.
- Jastremski, C. A., & Harvey, M. (1998). Making changes to improve the intensive care unit experience for patients and their families. *New Horizons, 6*(1), 99-109.
- Johnson, A. N. (2001). Neonatal response to control of noise inside the incubator. *Pediatric Nursing, 27*(6), 600-605.
- Johnson, A. N. (2003). Adapting the neonatal intensive care environment to decrease noise. *The Journal of Perinatal & Neonatal Nursing, 17*(4), 280-288; quiz 289-290.
- Kahn, D. M., Cook, T. E., Carlisle, C. C., Nelson, D. L., Kramer, N. R., & Millman, R. P. (1998). Identification and modification of environmental noise in an ICU setting. *Chest, 114*(2), 535-540.
- Kaldenburg, D. O. (1999). The influence of having a roommate on patient satisfaction. *Satisfaction Monitor*, January/February ([www.pressganey.org](http://www.pressganey.org)).
- Kam, P. C., Kam, A. C., & Thompson, J. F. (1994). Noise pollution in the anaesthetic and intensive care environment. *Anaesthesia, 49*(11), 982-986.
- Katcher, A., Segal, H., & Beck, A. (1984). Comparison of contemplation and hypnosis for the reduction of anxiety and discomfort during dental surgery. *American Journal of Clinical Hypnosis, 27*(1), 14-21.
- Kecskes, I., Rihmer, Z., Kiss, K., Vargha, A., Szili, I., & Rihmer, A. (2003). Possible effect of gender and season on the length of hospitalisation in unipolar major depressives. *Journal of Affective Disorders, 73*(3), 279-282.
- Keep, P. J. (1977). Stimulus deprivation in windowless rooms. *Anaesthesia, 32*(7), 598-602.
- Keep, P. J., James, J., & Inman, M. (1980). Windows in the intensive therapy unit. *Anaesthesia, 35*(3), 257-262.
- Keep, P. J. (1977). Stimulus deprivation in windowless rooms. *Anaesthesia, 32*(7), 598-602.
- Keipert, J. A. (1985). The harmful effects of noise in a children's ward. *Australian Paediatric Journal, 21*(2), 101-103.
- Kennedy, K., Fielder, A., Hardy, R., Tung, B., Gordon, D., & Reynolds, J. (2001). Reduced lighting does not improve medical outcomes in very low birth weight infants. *Journal of Pediatrics, 139*(4), 527-531.
- Kent, W. D., Tan, A. K., Clarke, M. C., & Bardell, T. (2002). Excessive noise levels in the neonatal ICU: Potential effects on auditory system development. *Journal of Otolaryngology, 31*(6), 355-360.
- Kinnunen, T., Saynajakangas, O., Tuuponen, T., & Keistinen, T. (2002). Regional and seasonal variation in the length of hospital stay for chronic obstructive pulmonary disease in Finland. *International Journal of Circumpolar Health, 61*(2), 131-135.
- Kulik, J. A., & Mahler, H. I. (1987). Effects of preoperative roommate assignment on preoperative anxiety and recovery from coronary-bypass surgery. *Health Psychology, 6*(6), 525-543.
- Kulik, J. A., Mahler, H. I., & Moore, P. J. (1996). Social comparison and affiliation under threat: Effects on recovery from major surgery. *Journal of Personality and Social Psychology, 71*(5), 967-979.
- Kulik, J. A., Moore, P. J., & Mahler, H. I. (1993). Stress and affiliation: Hospital roommate effects on preoperative anxiety and social interaction. *Health Psychology, 12*(2), 118-124.
- Kuller, R., & Laike, T. (1998). The impact of flicker from fluorescent lighting on well-being, performance, and physiological arousal. *Ergonomics, 41*(4), 433-447.
- Kumar, S., & Ng, B. (2001). Crowding and violence on psychiatric wards: Explanatory models. *Canadian Journal of Psychiatry, 46*(5), 433-437.
- Lawn, R. (1976). *Direction finding in an hospital environment: Waikate Hospital, Hamilton, New Zealand*. Los Angeles: School of Architecture and Urban Planning, University of

- California at Los Angeles.
- Lawson, B. R., & Phiri, M. (2003). *The architectural healthcare environment and its effects on patient health outcomes: A report on an NHS estates funded research project*. London: The Stationery Office.
- Leather, P., Beale, D., Santos, A., Watts, J., & Lee, L. (2003). Outcomes of environmental appraisal of different hospital waiting areas. *Environment & Behavior*, 35(6), 842-869.
- Leather, P., Pyrgas, M., Beale, D., & Lawrence, C. (1998). Windows in the workplace: Sunlight, view, and occupational stress. *Environment & Behavior*, 30(6), 739-762.
- Lee, D. W., Chan, A. C., Wong, S. K., Fung, T. M., Li, A. C., Chan, S. K., et al. (2004). Can visual distraction decrease the dose of patient-controlled sedation required during colonoscopy? A prospective randomized controlled trial. *Endoscopy*, 36(3), 197-201.
- Levine, M., Marchon, I., & Hanley, G. (1984). The placement and misplacement of you-are-here maps. *Environment & Behavior*, 16(2), 139-157.
- Levy, G. D., Woolston, D. J., & Browne, J. V. (2003). Mean noise amounts in level II vs level III neonatal intensive care units. *Neonatal Network*, 22(2), 33-38.
- Lewis, M., Bendersky, M., Koons, A., Hegyi, T., Hiatt, I. M., Ostfeld, B., et al. (1991). Visitation to a neonatal intensive care unit. *Pediatrics*, 88(4), 795-800.
- Lewis, P., Staniland, J., Cuppage, A., & Davies, J. M. (1990). Operating room noise. *Canadian Journal of Anaesthesia*, 37(Pt 2), S79.
- Lewy, A. J., Bauer, V. K., Cutler, N. L., Sack, R. L., Ahmed, S., Thomas, K. H., et al. (1998). Morning vs evening light treatment of patients with winter depression. *Archives of General Psychiatry*, 55(10), 890-896.
- Lewy, A. J., Nurnberger, J. I., Wehr, T. A., Pack, D., Becker, L. E., Powell R., et al. (1985). Supersensitivity to light: Possible trait marker for manic-depressive illness. *American Journal of Psychiatry*, 142(6), 725-727.
- Lingard, L., Reznick, R., Espin, S., Regehr, G., & DeVito, I. (2002). Team communications in the operating room: Talk patterns, sites of tension, and implications for novices. *Academic Medicine*, 77(3), 232-237.
- Liu, E. H., & Tan, S. (2000). Patients' perception of sound levels in the surgical suite. *Journal of Clinical Anesthesia*, 12(4), 298-302.
- Lopez, J., & Hendrickson, S. (1991). Family visits and different cultures. *Axone*, 12(3), 59-62.
- Lotas, M. J. (1992). Effects of light and sound in the neonatal intensive care unit environment on the low-birth-weight infant. *NAACOG's Clinical Issues in Perinatal and Women's Health Nursing*, 3(1), 34-44.
- Love, H. (2003). Noise exposure in the orthopaedic operating theatre: A significant health hazard. *ANZ Journal of Surgery*, 73(10), 836-838.
- Lovell, B. B., Ancoli-Israel, S., & Gevirtz, R. (1995). Effect of bright light treatment on agitated behavior in institutionalized elderly subjects. *Psychiatry Research*, 57(1), 7-12.
- Madi, B. C., Sandall, J., Bennett, R., & MacLeod, C. (1999). Effects of female relative support in labor: A randomized controlled trial. *Birth*, 26(1), 4-8.
- Malkin, J. (1989). Wayfinding: An orientation system for hospitals. *Progressive Architecture*, 70(12), 107-108.
- Mann, N., Haddow, R., Stokes, L., Goodley, S., & Rutter, N. (1986). Effect of night and day on preterm infants in a newborn nursery. *British Medical Journal*, 293(6557), 1265-1267.
- McCarthy, D. O., Ouimet, M. E., & Daun, J. M. (1991). Shades of Florence Nightingale: Potential impact of noise stress on wound healing. *Holistic Nursing Practice*, 5(4), 39-48.
- McGonagle, I. M., & Allan, S. (2002). A comparison of behaviour in two differing psychiatric long-stay rehabilitation environments. *Journal of Psychiatric and Mental Health Nursing*, 9(4), 493-499.
- McIntosh, S., Schuessler, B., Coron, D., Suchetka, M., Westball, C., & Dayton, D. (1994). What,

- if anything, does your unit do to reduce noise levels in the ICU? *Critical Care Nurse*, 14(6), 88-89.
- McLaughlin, A., McLaughlin, B., Elliott, J., & Campalani, G. (1996). Noise levels in a cardiac surgical intensive care unit: A preliminary study conducted in secret. *Intensive & Critical Care Nursing*, 12(4), 226-230.
- McLaughlin, J., Beebe, J., Hirshfield, P., Lindia, P., & Gubanc, D. (1996). *Duke University's Bird Garden*. Paper presented at the 1996 Annual Conference of the Society for the Arts in Healthcare, Durham, NC: Durham Arts Council and Duke University Medical Center.
- Mead, C. A. (1992). *Design issues related to hospital wayfinding*. Unpublished master's thesis, California State University, Long Beach.
- Meyer, T. J., Eveloff, S. E., Bauer, M. S., Schwartz, W. A., Hill, N. S., & Millman, R. P. (1994). Adverse environmental conditions in the respiratory and medical ICU settings. *Chest*, 105(4), 1211-1216.
- Meyer-Falcke, A., Rack, R., Eichwede, F., & Jansing, P. J. (1994). How noisy are anaesthesia and intensive care medicine? Quantification of the patients' stress. *European Journal of Anaesthesiology*, 11(5), 407-411.
- Miller, A. C., Hickman, L. C., & Lemasters, G. K. (1992). A distraction technique for control of burn pain. *Journal of Burn Care and Rehabilitation*, 13(5), 576-580.
- Miller, C., & Lewis, D. (1999). *Wayfinding: Effective wayfinding and signing systems: Guidance for healthcare facilities*. London: NHS Estates.
- Miller, C. L., White, R., Whitman, T. L., O'Callaghan, M. F., & Maxwell, S. E. (1995). The effects of cycled versus noncycled lighting on growth and development in preterm infants. *Infant Behavior and Development*, 18(1), 87-95.
- Minckley, B. B. (1968). A study of noise and its relationship to patient discomfort in the recovery room. *Nursing Research*, 17(3), 247-250.
- Moeser, S. D. (1988). Cognitive mapping in a complex building. *Environment & Behavior*, 20(1), 21-49.
- Montello, D. (1991). Spatial orientation and the angularity of urban routes: A field study. *Environment & Behavior*, 23(1), 47-69.
- Moore, M. M., Nguyen, D., Nolan, S. P., Robinson, S. P., Ryals, B., Imbrie, J. Z., et al. (1998). Interventions to reduce decibel levels on patient care units. *American Surgeon*, 64(9), 894.
- Morrison, W. E., Haas, E. C., Shaffner, D. H., Garrett, E. S., & Fackler, J. C. (2003). Noise, stress, and annoyance in a pediatric intensive care unit. *Critical Care Medicine*, 31(1), 113-119.
- Morse, J. M., & Pooler, C. (2002). Patient-family-nurse interactions in the trauma-resuscitation room. *American Journal of Critical Care*, 11(3), 240-249.
- Mullett, H., Synnott, K., & Quinlan, W. (1999). Occupational noise levels in orthopaedic surgery. *Irish Journal of Medical Science*, 168(2), 106.
- Murthy, V. S., Malhotra, S. K., Bala, I., & Raghunathan, M. (1995). Auditory functions in anaesthesia residents during exposure to operating room noise. *Indian Journal of Medical Research*, 101, 213-216.
- Nelson-Shulman, Y. (1983-84). Information and environmental stress: Report of a hospital intervention. *Journal of Environmental Systems*, 13(4), 303-316.
- Norbeck, J. S. (1985). Perceived job stress, job satisfaction, and psychological symptoms in critical care nursing. *Research in Nursing & Health*, 8(3), 253-259.
- Nott, M. R., & West, P. D. (2003). Orthopaedic theatre noise: A potential hazard to patients. *Anaesthesia*, 58(8), 784-787.
- Novaes, M. A., Aronovich, A., Ferraz, M. B., & Knobel, E. (1997). Stressors in ICU: Patients' evaluation. *Intensive Care Medicine*, 23(12), 1282-1285.

- Nystrom, K., & Axelsson, K. (2002). Mothers' experience of being separated from their newborns. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 31(3), 275-282.
- Nzama, N. P., Nolte, A. G., & Dorfling, C. S. (1995). Noise in a neonatal unit: Guidelines for the reduction or prevention of noise. *Curationis*, 18(2), 16-21.
- O'Connor, B. P., Davidson, H., & Gifford, R. (1991). Window view, social exposure and nursing home adaptation. *Canadian Journal on Aging*, 10(3), 216-223.
- Ogilvie, A. J. (1980). Sources and levels of noise on the ward at night. *Nursing Times*, 76(31), 1363-1366.
- Oren, D. A. (1999). There is nothing new under the sun. *American Journal of Psychiatry*, 156(2), 336.
- Oren, D. A., Brainard, G. C., Johnston, S. H., Joseph-Vanderpool, J. R., Sorek, E., & Rosenthal, N. E. (1991). Treatment of seasonal affective disorder with green light and red light. *American Journal of Psychiatry*, 148(4), 509.
- Oren, D. A., Cubells, J. F., & Litsch, S. (2001). Bright light for schizoaffective disorder. *American Journal of Psychiatry*, 158(12), 2086.
- Oren, D. A., Wisner, K. L., Spinelli, M., Epperson, C. N., Peindl, K. S., Terman, J. S., et al. (2002). An open trial of morning light therapy for treatment of antepartum depression. *American Journal of Psychiatry*, 159(4), 666.
- Ortega-Andeane, P. (1991). Physical comfort, social contact, wayfinding and meaning as indicators of stress in a hospital environment. *Proceedings of the Environmental Design Research Association, USA*, 22, 41-47.
- Ortega-Andeane, P., & Urbina-Soria, J. (1988). A case study of wayfinding and security in a Mexico City hospital. *Proceedings of the Environmental Design Research Association, USA*, 19, 231-236.
- Parsons, R., & Hartig, T. (2000). Environmental psychophysiology. In J. T. Cacioppo & L. G. Tassinary (Eds.), *Handbook of psychophysiology (2nd ed.)* (pp. 815-846). New York: Cambridge University Press.
- Parthasarathy, S., & Tobin, M. J. (2004). Sleep in the intensive care unit. *Intensive Care Medicine*, 30(2), 197-206.
- Partonen, T., & Lonnqvist, J. (2000). Bright light improves vitality and alleviates distress in healthy people. *Journal of Affective Disorders*, 57(1-3), 55-61.
- Passini, R., Rainville, C., Marchand, N., & Joannette, Y. (1995). Wayfinding in dementia of the Alzheimer type: Planning abilities. *Journal of Clinical and Experimental Neuropsychology* 17(6), 820-832.
- Pattison, H. M., & Robertson, C. E. (1996). The effect of ward design on the well-being of post-operative patients. *Journal of Advanced Nursing*, 23(4), 820-826.
- Peponis, J., Zimring, C., & Choi, Y. K. (1990). Finding the building in wayfinding. *Environment & Behavior*, 22(5), 555-590.
- Peterson, R., Knapp, T., Rosen, J., & al., e. (1977). The effects of furniture arrangement on the behavior of geriatric patients. *Behavior Therapy*, 8, 464-467.
- Persson Waye, K., Rylander, R., Benton, S., & Leventhall, H. G. (1997). Effects on performance and work quality due to low frequency ventilation noise. *Journal of Sound and Vibration*, 205(4), 467-474.
- Peterson, R., Knapp, T., Rosen, J., & Pither, B. F. (1977). The effects of furniture arrangement on the behavior of geriatric patients. *Behavior Therapy*, 8(3), 464-467.
- Philbin, M. K., & Gray, L. (2002). Changing levels of quiet in an intensive care nursery. *Journal of Perinatology*, 22(6), 455-460.
- Port, C. L., Hebel, J. R., Gruber-Baldini, A. L., Baumgarten, M., Burton, L., Zimmerman, S., et al. (2003). Measuring the frequency of contact between nursing home residents and their family and friends. *Nursing Research*, 52(1), 52-56.

- Powell, M. (1973). Building noise in a hospital: An experimental simulation. *Annals of Occupational Hygiene*, 16(1), 77-79.
- Powers, K. S., & Rubenstein, J. S. (1999). Family presence during invasive procedures in the pediatric intensive care unit: A prospective study. *Archives of Pediatrics & Adolescent Medicine*, 153(9), 955-958.
- Pragay, D. A. (1981). Noise levels in hospital laboratories. Are they a problem and can they be reduced? *Clinical Biochemistry*, 14(3), 157-158.
- Raeside, L. (1997). Perceptions of environmental stressors in the neonatal unit. *British Journal of Nursing*, 6(16), 914-916, 918, 920-913.
- Rainville, C., Passini, R., & Marchand, N. (2001). A multiple case study of wayfinding in dementia of the Alzheimer type: Decision making. *Aging, Neuropsychology, & Cognition*, 8(1), 54.
- Ray, C. D., & Levinson, R. (1992). Noise pollution in the operating room: A hazard to surgeons, personnel, and patients. *Journal of Spinal Disorders*, 5(4), 485-488.
- Redding, J. S., Hargest, T. S., & Minsky, S. H. (1977). How noisy is intensive care? *Critical Care Medicine*, 5(6), 275-276.
- Regan, T. (1971). *Wayfinding in hospital environments, an overview*. Los Angeles: School of Architecture and Urban Planning, University of California at Los Angeles.
- Rice, C. G., Talbott, J. A., & Stern, D. (1980). Effects of environmental agents on social behavior of patients in a hospital dining room. *Hospital & Community Psychiatry*, 31(2), 128-130.
- Robertson, A., Cooper-Peel, C., & Vos, P. (1998). Peak noise distribution in the neonatal intensive care nursery. *Journal of Perinatology*, 18(5), 361-364.
- Robertson, A., Cooper-Peel, C., & Vos, P. (1999). Contribution of heating, ventilation, and air conditioning airflow and conversation to the ambient sound in a neonatal intensive care unit. *Journal of Perinatology*, 19(5), 362-366.
- Rosenfield, A. G. (1980). Visiting in the intensive care nursery. *Child Development*, 51(3), 939-941.
- Rosenthal, N. E., Sack, D. A., Carpenter, C. J., Parry, B. L., Mendelson, W. B., & Wehr, T. A. (1985). Antidepressant effects of light in seasonal affective disorder. *American Journal of Psychiatry*, 142(2), 163-170.
- Routhieaux, R. L., & Tansik, D. A. (1997). The benefits of music in hospital waiting rooms. *Health Care Supervisor*, 16(2), 31-40.
- Sadeh, A., McGuire, J. P., Sachs, H., Seifer, R., Tremblay, A., Civita, R., et al. (1995). Sleep and psychological characteristics of children on a psychiatric inpatient unit. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34(6), 813-819.
- Sallstrom, C., Sandman, P. O., & Norberg, A. (1987). Relatives' experience of the terminal care of long-term geriatric patients in open-plan rooms. *Scandinavian Journal of Caring Sciences*, 1(4), 133-140.
- Satlin, A., Volicer, L., Ross, V., Herz, L., & Campbell, S. (1992). Bright light treatment of behavioral and sleep disturbances. *American Journal of Psychiatry*, 149(8), 1028.
- Saunders, A. N. (1995). Incubator noise: A method to decrease decibels. *Pediatric Nursing*, 21(3), 265-268.
- Schneider, L. F., & Taylor, H. A. (1999). How do you get there from here? Mental representations of route descriptions. *Applied Cognitive Psychology*, 13(5), 415-441.
- Schneider, S. M., Ellis, M., Coombs, W. T., Shonkwiler, E. L., & Folsom, L. C. (2003). Virtual reality intervention for older women with breast cancer. *Cyberpsychology & Behavior*, 6(3), 301-307.
- Schneider, S. M., Gallery, M. E., Schafermeyer, R., & Zwemer, F. L. (2003). Emergency department crowding: A point in time. *Annals of Emergency Medicine*, 42(2), 167-172.
- Schneider, S. M., Prince-Paul, M., Allen, M. J., Silverman, P., & Talaba, D. (2004). Virtual

- reality as a distraction intervention for women receiving chemotherapy. *Oncology Nursing Forum*, 31(1), 81-88.
- Schnelle, J. F., Cruise, P. A., Alessi, C. A., Ludlow, K., al-Samarrai, N. R., & Ouslander, J. G. (1998). Sleep hygiene in physically dependent nursing home residents: Behavioral and environmental intervention implications. *Sleep*, 21(5), 515-523.
- Schnelle, J. F., Ouslander, J. G., Simmons, S. F., Alessi, C. A., & Gravel, M. D. (1993). The nighttime environment, incontinence care, and sleep disruption in nursing homes. *Journal of the American Geriatrics Society*, 41(9), 910-914.
- Schofield, P., & Davis, B. (2000). Sensory stimulation (snoezelen) versus relaxation: A potential strategy for the management of chronic pain. *Disability & Rehabilitation*, 22(15), 675-682.
- Schulte, D. A., Burrell, L. O., Gueldner, S. H., Bramlett, M. H., Fuszard, B., Stone, S. K., et al. (1993). Pilot study of the relationship between heart rate and ectopy and unrestricted vs restricted visiting hours in the coronary care unit. *American Journal of Critical Care*, 2(2), 134-136.
- Schuster, R. J., & Weber, M. L. (2003). Noise in the ambulatory health care setting. How loud is too loud? *Journal of Ambulatory Care Management*, 26(3), 243-249.
- Scott, R. A. (1995). Beeper technology: Improving staff efficiency and decreasing noise levels on medical-surgical units. *Medsurg Nursing*, 4(1), 73-75.
- Shankar, N., Malhotra, K. L., Ahuja, S., & Tandon, O. P. (2001). Noise pollution: A study of noise levels in the operation theatres of a general hospital during various surgical procedures. *Journal of the Indian Medical Association*, 99(5), 244, 246-247.
- Shapiro, R. A., & Berland, T. (1972). Noise in the operating room. *New England Journal of Medicine*, 287(24), 1236-1238.
- Shertzer, K. E., & Keck, J. F. (2001). Music and the PACU environment. *Journal of Perianesthesia Nursing*, 16(2), 90-102.
- Shogan, M. G., & Schumann, L. L. (1993). The effect of environmental lighting on the oxygen saturation of preterm infants in the NICU. *Neonatal Network*, 12(5), 7-13.
- Simons, J. W., Mize, J. J., & Haynes, B. C., Jr. (1969). Acoustical properties of carpets and drapes. *Medical Journal of Australia*, 1(17), 125-127.
- Simpson, T. (1992). Visitors' verbal behavior with coronary care unit patients. *Western Journal of Nursing Research*, 14(4), 482-494; discussion 495-487.
- Simpson, T. (1993). Visit preferences of middle-aged vs older critically ill patients. *American Journal of Critical Care*, 2(4), 339-345.
- Simpson, T., Lee, E. R., & Cameron, C. (1996). Patients' perceptions of environmental factors that disturb sleep after cardiac surgery. *American Journal of Critical Care*, 5(3), 173-181.
- Simpson, T., Lee, E. R., & Cameron, C. (1996). Relationships among sleep dimensions and factors that impair sleep after cardiac surgery. *Research in Nursing & Health*, 19(3), 213-223.
- Simpson, T., & Shaver, J. (1991). A comparison of hypertensive and nonhypertensive coronary care patients' cardiovascular responses to visitors. *Heart & Lung*, 20(3), 213-220.
- Slevin, M., Farrington, N., Duffy, G., Daly, L., & Murphy, J. F. (2000). Altering the NICU and measuring infants' responses. *Acta Paediatrica*, 89(5), 577-581.
- Snyder-Halpern, R. (1985). The effect of critical care unit noise on patient sleep cycles. *Critical Care Quarterly*, 7(4), 41-51.
- Sommer, R., & Ross, H. (1958). Social interaction on a geriatrics ward. *International Journal of Social Psychiatry*, 4(2), 128-133.
- Soutar, R. L., & Wilson, J. A. (1986). Does hospital noise disturb patients? *British Medical Journal (Clinical Research Ed.)*, 292(6516), 305.
- Southwell, M. T., & Wistow, G. (1995). Sleep in hospitals at night: Are patients' needs being

- met? *Journal of Advanced Nursing*, 21(6), 1101-1109.
- Stanton, N., & Edworthy, J. (1998). Auditory affordances in the intensive treatment unit. *Applied Ergonomics*, 29(5), 389-394.
- Starks, M. A. (2003). Restoring attention in pregnancy: The natural environment. *Clinical Nursing Research*, 12(3), 246-265.
- Stoneham, J., & Jones, R. (1997). Residential landscapes: Their contribution to the quality of older people's lives. *Activities, Adaptation & Aging*, 22(1), 17-26.
- Storch, D. D. (2001). Climate and length of hospital stay. *American Journal of Psychiatry*, 158(9), 1530.
- Strauch, C., Brandt, S., & Edwards-Beckett, J. (1993). Implementation of a quiet hour: Effect on noise levels and infant sleep states. *Neonatal Network* 12(2), 31-35.
- Stroud, R. (1997). The effects of technology on relatives in critical care environments. *Nursing in Critical Care*, 2(6), 272-275.
- Sullivan, E. E. (2001). Family visitation in PACU. *Journal of Perianesthesia Nursing*, 16(1), 29-30.
- Sumaya, I. C., Rienzi, B. M., Deegan, II, J. F., & Moss, D. E. (2001). Bright light treatment decreases depression in institutionalized older adults: A placebo-controlled crossover study. *Journal of Gerontology*, 56A(6), M356-M360.
- Surethiran, S. S., Wilbraham, K., May, J., Chant, T., Emmerson, A. J., & Newton, V. E. (2003). Noise levels within the ear and post-nasal space in neonates in intensive care. *Archives of Disease in Childhood. Fetal and Neonatal Edition*, 88(4), F315-318.
- Tanimoto, S., Takayanagi, K., Yokota, H., & Yamamoto, Y. (1999). The psychological and physiological effects of an intensive-care unit environment on healthy individuals. *Clinical Performance and Quality Health Care*, 7(2), 77-82.
- Terman, J. S., Terman, M., Lo, E. S., & Cooper, T. B. (2001). Circadian time of morning light administration and therapeutic response in winter depression. *Archives of General Psychiatry*, 58(1), 69-75.
- Terman, M., Terman, J. S., & Ross, D. C. (1998). A controlled trial of timed bright light and negative air ionization for treatment of winter depression. *Archives of General Psychiatry*, 55(10), 875.
- Thomas, K. A. (1990). Design issues in the NICU: Thermal effects of windows. *Neonatal Network*, 9(4), 23-26.
- Thomas, K. A., & Martin, P. A. (2000). NICU sound environment and the potential problems for caregivers. *Journal of Perinatology*, 20(Pt 2), S94-99.
- Thompson, T., Robinson, J., Dietrich, M., Farris, M., & Sinclair, V. (1996). Architectural features and perceptions of community residences for people with mental retardation. *American Journal of Mental Retardation*, 101(3), 292-314.
- Thornes, R. (1983). Parental access and family facilities in children's wards in England. *British Medical Journal (Clinical Research Ed.)*, 287(6386), 190-192.
- Topf, M. (1983). Noise pollution in the hospital. *New England Journal of Medicine*, 309(1), 53-54.
- Topf, M. (1985). Noise-induced stress in hospital patients: Coping and nonauditory health outcomes. *Journal of Human Stress*, 11(3), 125-134.
- Topf, M. (1985). Personal and environmental predictors of patient disturbance due to hospital noise. *Journal of Applied Psychology*, 70(1), 22-28.
- Topf, M. (1988). Noise-induced occupational stress and health in critical care nurses. *Hospital Topics*, 66(1), 30-34.
- Topf, M. (1989). Sensitivity to noise, personality hardiness, and noise-induced stress in critical care nurses. *Environment and Behavior*, 21(6), 717-733.
- Topf, M. (1992). Stress effects of personal control over hospital noise. *Behavioral Medicine*,

- 18(2), 84-94.
- Topf, M. (1992). Effects of personal control over hospital noise on sleep. *Research in Nursing & Health, 15*(1), 19-28.
- Topf, M. (1999). Does the environment of the hospital affect recovery? *Journal of Health Services Research & Policy, 4*(2), 127.
- Topf, M., Bookman, M., & Arand, D. (1996). Effects of critical care unit noise on the subjective quality of sleep. *Journal of Advanced Nursing, 24*(3), 545-551.
- Topf, M., & Davis, J. E. (1993). Critical care unit noise and rapid eye movement (REM) sleep. *Heart & Lung, 22*(3), 252-258.
- Topf, M., & Dillon, E. (1988). Noise-induced stress as a predictor of burnout in critical care nurses. *Heart & Lung, 17*(5), 567-574.
- Topf, M., & Thompson, S. (2001). Interactive relationships between hospital patients' noise-induced stress and other stress with sleep. *Heart & Lung, 30*(4), 237-243.
- Trites, D. K., & Green, R. M. (1970). Hospital visiting: The patients' point of view. *Nursing Outlook, 18*(8), 44-45.
- Tse, M. M. Y., Ng, J. K. F., Chung, J. W. Y., & Wong, T. K. S. (2002). The effect of visual stimuli on pain threshold and tolerance. *Journal of Clinical Nursing, 11*(4), 462-469.
- Tse, M. M. Y., Ng, J. K. F., Chung, J. W. Y., & Wong, T. K. S. (2002). The effect of visual stimulation via the eyeglass display and the perception of pain. *CyberPsychology & Behavior, 5*(1), 65-75.
- Tsiou, C., Eftymiatis, D., Theodossopoulou, E., Notis, P., & Kiriakou, K. (1998). Noise sources and levels in the Evgenidion Hospital intensive care unit. *Intensive Care Medicine, 24*(8), 845-847.
- Ulrich, R. S. (1984). View through a window may influence recovery from surgery. *Science, 224*(4647), 420-421.
- Ulrich, R. S. (1991). Effects of interior design on wellness: Theory and recent scientific research. *Journal of Health Care Interior Design, 3*(1), 97-109.
- Ulrich, R. S. (1999). Effects of gardens on health outcomes: Theory and research. In C. Cooper Marcus & M. Barnes (Eds.), *Healing Gardens* (pp. 27-86). New York: Wiley.
- Ulrich, R. S., & Gilpin, L. (2003). Healing arts: Nutrition for the soul. In S. B. Frampton, L. Gilpin & P. Charmel (Eds.), *Putting patients first: Designing and practicing patient-centered care* (pp. 117-146). San Francisco: Jossey-Bass.
- Ulrich, R. S., Lawson, B., & Martinez, M. (2003). *Exploring the patient environment: An NHS estates workshop*. London: The Stationery Office.
- Ulrich, R. S., Lunden, O., & Eltinge, J. L. (1993). *Effects of exposure to nature and abstract pictures on patients recovering from heart surgery*. Paper presented at the thirty-third meeting of the Society for Psychophysiological Research, Rottach-Egern, Germany. Abstract in *Psychophysiology, 30*(Suppl. 1), 7.
- Ulrich, R. S., Simons, R. F., Losito, B. D., Fiorito, E., Miles, M. A., & Zelson, M. (1991). Stress recovery during exposure to natural and urban environments. *Journal of Environmental Psychology, 11*(2): 201-230.
- Ulrich, R. S., Simons, R. F., & Miles, M. A. (2003). Effects of environmental simulations and television on blood donor stress. *Journal of Architectural & Planning Research, 20*(1), 38-47.
- Van der Ploeg, H. M. (1988). Stressful medical events: A survey of patients' perceptions. In S. Maies, C. D. Spielberger, P. B. Defares & I. G. Sarason (Eds.), *Topics in Health Psychology* (pp. 193-203). New York: John Wiley.
- Van Someren, E. J. W., Kessler, A., Mirmiran, M., & Swaab, D. F. (1997). Indirect bright light improves circadian rest-activity rhythm disturbances in demented patients. *Biological Psychiatry, 41*(9), 955-963.



- Verderber, S. (1986). Dimensions of person-window transactions in the hospital environment. *Environment & Behavior*, 18(4), 450-466.
- Verderber, S., & Reuman, D. (1987). Windows, views, and health status in hospital therapeutic environments. *Journal of Architectural & Planning Research*, 4(2), 120-133.
- Vessey, J. A., Carlson, K. L., & McGill, J. (1994). Use of distraction with children during an acute pain experience. *Nursing Research*, 43(6), 369-372.
- Vickery, K. (2002). Study views visiting patterns. *Provider*, 28(11), 45-46, 49.
- Vogelsang, J. (1988). Effect of visitors on patient behavior in the postanesthesia period. *Dimensions of Critical Care Nursing*, 7(2), 91-100.
- Volicer, B. J., Isenberg, M. A., & Burns, M. W. (1977). Medical-surgical differences in hospital stress factors. *Journal of Human Stress*, 3(2), 3-13.
- Walch, J. M., Rabin, B. S., Day, R., Williams, J. N., Choi, K., & Kang, J. D. (In press). The effect of sunlight on post-operative analgesic medication usage: A prospective study of spinal surgery patients.
- Walder, B., Francioli, D., Meyer, J. J., Lancon, M., & Romand, J. A. (2000). Effects of guidelines implementation in a surgical intensive care unit to control nighttime light and noise levels. *Critical Care Medicine*, 28(7), 2242-2247.
- Walker, J. S., Eakes, G. G., & Siebelink, E. (1998). The effects of familial voice interventions on comatose head-injured patients. *Journal of Trauma Nursing*, 5(2), 41-45.
- Wallace, C. J., Robins, J., Alvord, L. S., & Walker, J. M. (1999). The effect of earplugs on sleep measures during exposure to simulated intensive care unit noise. *American Journal of Critical Care*, 8(4), 210-219.
- Wallace-Guy, G., Kripke, D., Jean-Louis, G., Langer, R., Elliott, J., & Tuunainen, A. (2002). Evening light exposure: Implications for sleep and depression. *Journal of the American Geriatrics Society*, 50(4), 738-739.
- Walsh-Sukys, M., Reitenbach, A., Hudson-Barr, D., & DePompei, P. (2001). Reducing light and sound in the neonatal intensive care unit: An evaluation of patient safety, staff satisfaction and costs. *Journal of Perinatology*, 21(4), 230-235.
- Warren, N. A. (1993). Perceived needs of the family members in the critical care waiting room. *Critical Care Nursing Quarterly*, 16(3), 56-63.
- Weisman, J. (1981). Evaluating architectural legibility: Wayfinding in the built environment. *Environment & Behavior*, 13(2), 189-204.
- Whalen, L. (1992). Noise in the intensive care setting. *Canadian Critical Care Nursing Journal*, 8(1), 9-10.
- Whall, A. L., Black, M. E., Groh, C. J., Yankou, D. J., Kupferschmid, B. J., & Foster, N. L. (1997). The effect of natural environments upon agitation and aggression in late stage dementia patients. *American Journal of Alzheimer's Disease and Other Dementias*, September-October: 216-220.
- White, A., & Burgess, M. (1992). Strategies for reduction of noise levels in ICUs. *Australian Journal of Advanced Nursing*, 10(2), 22-26.
- White, R. D. (2003). Individual rooms in the NICU—an evolving concept. *Journal of Perinatology*, 23(1), S22-24.
- Whitehouse, S., Varni, J. W., Seid, M., Cooper-Marcus, C., Ensberg, M. J., Jacobs, J. R., et al. (2001). Evaluating a children's hospital garden environment: Utilization and consumer satisfaction. *Journal of Environmental Psychology*, 21(3), 301-314.
- Whitis, G. (1994). Visiting hospitalized patients. *Journal of Advanced Nursing*, 19(1), 85-88.
- Williams, M. A. (1988). The physical environment and patient care. *Annual Review of Nursing Research*, 6, 61-84. Williams, M. A. (1989). Physical environment of the intensive care unit and elderly patients. *Critical Care Nursing Quarterly*, 12(1), 52-60.
- Wilson, L. M. (1972). Intensive care delirium: The effect of outside deprivation in a windowless

- unit. *Archives of Internal Medicine*, 130(2), 225-226.
- Wirz-Justice, A. (1998). Beginning to see the light. *Archives of General Psychiatry*, 55(10), 861.
- Wood, A. M. (1993). A review of literature relating to sleep in hospital with emphasis on the sleep of the ICU patient. *Intensive & Critical Care Nursing*, 9(2), 129-136.
- Wright, P., Hull, A. J., & Lickorish, A. (1993). Navigating in a hospital outpatients' department: The merits of maps and wallsigns. *Journal of Architectural and Planning Research*, 10(1), 76-89.
- Wysocki, A. B. (1996). The effect of intermittent noise exposure on wound healing. *Advances in Wound Care*, 9(1), 35-39.
- Yamada, N., Martin-Iverson, M. T., Daimon, K., Tsujimoto, T., & Takahashi, S. (1995). Clinical and chronobiological effects of light therapy on nonseasonal affective disorders. *Biological Psychiatry*, 37(12), 866-873.
- Yamamoto-Mitani, N., Aneshensel, C. S., & Levy-Storms, L. (2002). Patterns of family visiting with institutionalized elders: The case of dementia. *Journals of Gerontology. Series B, Psychological Sciences and Social Sciences*, 57(4), S234-246.
- Yarcheski, A., & Knapp-Spooner, C. (1994). Stressors associated with coronary bypass surgery. *Clinical Nursing Research*, 3(1), 57-68.
- Yassi, A., Gaborieau, D., Gillespie, I., & Elias, J. (1991). The noise hazard in a large health care facility. *Journal of Occupational Medicine*, 33(10), 1067-1070.
- Yinon, A. M., Ilan, Y., Tadmor, B., Altarescu, G., & Hershko, C. (1992). Quality of sleep in the medical department. *British Journal of Clinical Practice*, 46(2), 88-91.
- Youngner, S. J., Coulton, C., Welton, R., Juknialis, B., & Jackson, D. L. (1984). ICU visiting policies. *Critical Care Medicine*, 12(7), 606-608.
- Zahr, L. K., & de Traversay, J. (1995). Premature infant responses to noise reduction by earmuffs: Effects on behavioral and physiologic measures. *Journal of Perinatology*, 15(6), 448-455.
- Zimring, C. (1990). *The costs of confusion: Non-monetary and monetary costs of the Emory University hospital wayfinding system*. Atlanta: Georgia Institute of Technology.
- Zimring, C., & Templer, J. (1983-84). Wayfinding and orientation by the visually impaired. *Journal of Environmental Systems*, 13(4), 333-352.

### ***Improve Overall Health-Care Quality***

- Arneill, A. B., & Devlin, A. S. (2002). Perceived quality of care: The influence of the waiting room environment. *Journal of Environmental Psychology*, 22(4), 345-360.
- Bostick, R., & Krebs, M. (1983). The spinal cord injured female in the veterans hospital: The role of the environment in rehabilitation. *Journal of the American Paraplegia Society*, 6(4), 81-84.
- Boswell, K., Finlay, F., Jones, R., & Hill, P. (2000). Perceived ideal out-patient department and hospital ward for children, adolescents and their families. *Clinical Child Psychology & Psychiatry*, 5(2), 213-219.
- Bruster, S., Jarman, B., Bosanquet, N., Weston, D., Erens, R., & Delbanco, T. L. (1994). National survey of hospital patients. *British Medical Journal (Clinical Research Ed.)*, 309(6968), 1542-1546.
- Caplan, G., Board, N., Paten, A., Tazelaar-Molinia, J., Crowe, P., Yap, S. J., et al. (1999). Decreasing lengths of stay: The cost to the community. *The Australian and New Zealand Journal of Surgery*, 69(6), 433-437.
- Chaudhury, H., Mahmood, A., & Valente, M. (2003). *Pilot study on comparative assessment of patient care issues in single and multiple occupancy rooms*. Unpublished report, The

- Coalition for Health Environments Research.
- Chen, T. S., & Sanoff, H. (1988). The patients' view of their domain. *Design Studies*, 9(1), 40-55.
- Counsell, S. R., Holder, C. M., Liebenauer, L. L., Palmer, R. M., Fortinsky, R. H., Kresevic, D. M., et al. (2000). Effects of a multicomponent intervention on functional outcomes and process of care in hospitalized older patients: A randomized controlled trial of Acute Care for Elders (ACE) in a community hospital. *Journal of the American Geriatrics Society*, 48(12), 1572-1581.
- Devlin, A. S. (1992). Psychiatric ward renovation: Staff perception and patient behavior. *Environment & Behavior*, 24(1), 66-84.
- Devlin, A. S. (1995). Staff, patients, and visitors: Responses to hospital unit enhancements. *Proceedings of the Environmental Design Research Association*, 26, 113-117.
- Devlin, A. S., & Arneill, A. B. (2003). Health care environments and patient outcomes: A review of the literature. *Environment & Behavior*, 35(5), 665-694.
- Dolce, J. J., Doleys, D. M., Raczynski, J. M., & Crocker, M. F. (1985). Narcotic utilization for back pain patients housed in private and semi-private rooms. *Addictive Behaviors*, 10(1), 91-95.
- Fanurik, D., Schmitz, M. L., Martin, G. A., Koh, J. L., Wood, M., Sturgeon, L., et al. (2000). Hospital room or treatment room: Where should inpatient pediatric procedures be performed? *Children's Health Care*, 29(2), 103-111.
- Fowler, E., MacRae, S., Stern, A., Harrison, T., Gerteis, M., Walker, J., et al. (1999). The built environment as a component of quality care: Understanding and including the patient's perspective. *The Joint Commission Journal on Quality Improvement*, 25(7), 352-362.
- Francis, J. J., Pankratz, V. S., & Huddleston, J. M. (2001). Patient satisfaction associated with correct identification of physician's photographs. *Mayo Clinic Proceedings*, 76(6), 604-608.
- Gotlieb, J. B. (2000). Understanding the effects of nurses, patients' hospital rooms, and patients' perception of control in the perceived quality of a hospital. *Health Marketing Quarterly*, 18(1/2), 1.
- Griebel, M. A. (1985). *Post occupancy evaluation of hospitals: General methodology and application to five departments*. Unpublished master's thesis, Washington University, St. Louis, MO.
- Hahn, J. E., Jones, M. R., & Waszkiewicz, M. (1995). Renovation of a semiprivate patient room. Bowman Center Geriatric Rehabilitation Unit. *The Nursing Clinics of North America*, 30(1), 97-115.
- Happ, B. A. (1993). The effect of point of care technology on the quality of patient care. *Proceedings of Annual Symposium on Computer Application in Medical Care*, 17, 183-187.
- Harwood, R. H., & Ebrahim, S. (1992). Long-term institutional residents: Does the environment affect outcomes? *Journal of the Royal College of Physicians of London*, 26(2), 134-138.
- Holahan, C., & Saegert, S. (1973). Behavioral and attitudinal effects of large-scale variation in the physical environment of psychiatric wards. *Journal of Abnormal Psychology*, 82(3), 454-462.
- Hutton, A. (2003). Activities in the adolescent ward environment. *Contemporary Nurse*, 14(3), 312-319.
- Janssen, P. A., Klein, M. C., Harris, S. J., Soolsma, J., & Seymour, L. C. (2000). Single room maternity care and client satisfaction. *Birth*, 27(4), 235-243.
- Johansson, P., Oleni, M., & Fridlund, B. (2002). Patient satisfaction with nursing care in the context of health care: A literature study. *Scandinavian Journal of Caring Sciences*, 16(4), 337-344.
- Judkins, S. (2003). Paediatric emergency department design: Does it affect staff, patient and

- community satisfaction? *Emergency Medicine*, 15(1), 63-67.
- Kincaid, C., & Peacock, J. R. (2003). The effect of a wall mural on decreasing four types of door-testing behaviors. *Journal of Applied Gerontology*, 22(1), 76-88.
- Kirk, S. (2003). Single or shared rooms for hospice patients? *Palliative Medicine*, 17(6), 560.
- Koss, E., & Gilmore, G. C. (1998). Environmental interventions and functional ability of AD patients. In B. Vellas, J. Fitten & G. Frisoni (Eds.), *Research and Practice in Alzheimer's Disease* (pp. 185-193). New York: Springer.
- Krueckeberg, H. F., & Hubbert, A. (1995). Attribute correlates of hospital outpatient satisfaction. *Journal of Ambulatory Care Marketing*, 6(1), 11-43.
- Landefeld, C. S., Palmer, R. M., Kresevic, D. M., Fortinsky, R. H., & Kowal, J. (1995). A randomized trial of care in a hospital medical unit especially designed to improve the functional outcomes of acutely ill older patients. *New England Journal of Medicine*, 332(20), 1338-1344.
- Lawson, B., & Phiri, M. (2000). Hospital design: Room for improvement. *Health Services Journal*, 110(5688), 24-26.
- Linhorst, D. M. (1991). The use of single room occupancy (SRO) housing as a residential alternative for persons with a chronic mental illness. *Community Mental Health Journal*, 27(2), 135-144.
- Lyons, D., El Sayed, O. A., & Matthew, V. M. (2001). New unit on old ground or general hospital—where do patients want inpatient treatment? *Irish Journal of Psychological Medicine*, 18(4), 129-131.
- Martin, D. P., Diehr, P., Conrad, D. A., Davis, J. H., Leickly, R., & Perrin, E. B. (1998). Randomized trial of a patient-centered hospital unit. *Patient Education and Counseling*, 34(2), 125-133.
- Martin, D. P., Diehr, P., Conrad, D. A., Hunt Davis, J., Leickly, R., & Perrin, E. B. (1998). Randomized trial of a patient-centered hospital unit. *Patient Education and Counseling*, 34(2), 125-133.
- Melin, L., & Gotestam, K. G. (1981). The effects of rearranging ward routines on communication and eating behaviors of psychogeriatric patients. *Journal of Applied Behavior Analysis*, 14(1), 47-51.
- Middleboe, T., Schjodt, T., Byrstring, K., & Gjerris, A. (2001). Ward atmosphere in acute psychiatric in-patient care: Patients' perceptions, ideals and satisfaction. *Acta Psychiatrica Scandinavica*, 103(3), 212-219.
- Miller, N. O., Friedman, S. B., & Coupey, S. M. (1998). Adolescent preferences for rooming during hospitalization. *Journal of Adolescent Health*, 23(2), 89-93.
- Morgan, D. G., & Stewart, N. J. (1998). Multiple occupancy versus private rooms on dementia care units. *Environment & Behavior*, 30(4), 487-503.
- Morgan, D. G., & Stewart, N. J. (1998). High versus low density special care units: Impact on the behavior of elderly residents with dementia. *Canadian Journal on Aging*, 17(2), 143-165.
- Mujahed, M. (1999). *Design impact on health care facilities performance: A post-occupancy evaluation of patient rooms*. Unpublished master's thesis, Arizona State University, Tempe.
- Muller, M. J., Schlässer, R., Kapp-Steen, G., Schanz, B., & Benkert, O. (2002). Patients' satisfaction with psychiatric treatment: Comparison between an open and a closed ward. *Psychiatric Quarterly*, 73(2), 93.
- Olusina, A. K., Ohaeri, J. U., & Olatawura, M. O. (2002). Patient and staff satisfaction with the quality of in-patient psychiatric care in a Nigerian general hospital. *Social Psychiatry & Psychiatric Epidemiology*, 37(6), 283-286.
- Pease, N. J., & Finlay, I. G. (2002). Do patients and their relatives prefer single cubicles or shared wards? *Palliative Medicine*, 16(5), 445-446.

- Philips, C. (1988). Single room maternity care for maximum cost-efficiency. *Perinatology/Neonatology*, 12(March/April), 22-31.
- Pichert, J. W., Federspiel, C. F., Hickson, G. B., Miller, C. S., Gauld-Jaeger, J., & Gray, C. L. (1999). Identifying medical center units with disproportionate shares of patient complaints. *The Joint Commission Journal on Quality Improvement*, 25(6), 288-299.
- Press Ganey, Inc. (2003). National patient satisfaction data for 2003. (Provided by Press Ganey, Inc. at the request of the authors of this research review.)
- Seltzer, E., Schulman, K. A., Brennan, P. J., & Lynn, L. A. (1993). Patient attitudes toward rooming with persons with HIV infection. *Journal of Family Practice*, 37(6), 564-568.
- Shepley, M. M. (1995). The location of behavioral incidents in a children's psychiatric facility. *Children's Environments*, 12(3), 352-361.
- Shepley, M. M. (2002). Predesign and postoccupancy analysis of staff behavior in a neonatal intensive care unit. *Children's Health Care*, 31(3), 237-253.
- Shepley, M. M., & Wilson, P. (1999). Designing for persons with AIDS: A post-occupancy study at the Bailey-Boushay House. *Journal of Architectural and Planning Research*, 16(1), 17-32.
- Spaeth, G. L., & Angell, M. F. (1968). Preference of ophthalmic patients for multi-bedded hospital rooms. *Archives of Ophthalmology*, 79(4), 362-365.
- Stahler, G. J., Frazer, D., & Rappaport, H. (1984). The evaluation of an environmental remodeling program on a psychiatric geriatric ward. *Journal of Social Psychology*, 123(1), 101.
- Stegenga, P. W. (1990). Postsurgical recovery care: Spatial organization and social relationships. Unpublished master's thesis, Georgia Institute of Technology, Atlanta.
- Strittmatter, S., & Jessee, P. O. (1999). Pediatric health care: Parents' views on child life services. *Journal of Family and Consumer Sciences*, 91(1), 65.
- Sturdavant, M. (1960). Intensive nursing service in circular and rectangular units. *Hospitals, JAHA*, 34(14), 46-48, 71-78.
- Swan, J. E., Richardson, L. D., & Hutton, J. D. (2003). Do appealing hospital rooms increase patient evaluations of physicians, nurses, and hospital services? *Health Care Management Review*, 28(3), 254.
- Teresi, J. A., Holmes, D., & Monaco, C. (1993). An evaluation of the effects of commingling cognitively and noncognitively impaired individuals in long-term care facilities. *Gerontologist*, 33(3), 350-358.
- Thi, P. L. N., Briancon, S., Empereur, F., & Guillemin, F. (2002). Factors determining inpatient satisfaction with care. *Social Science & Medicine*, 54(4), 493-504.
- Watson, W. T., Marshall, E. S., & Fosbinder, D. (1999). Elderly patients' perceptions of care in the emergency department. *Journal of Emergency Nursing*, 25(2), 88-92.
- Willmott, M. (1986). The effect of a vinyl floor surface and a carpeted floor surface upon walking in elderly hospital in-patients. *Age and Ageing*, 15(2), 119-120.
- Winkelstein, M. L., & Carson, V. J. (1987). Adolescents and rooming-in. *Maternal-Child Nursing Journal*, 16(1), 75-88.